

See label for use on lawns, foundations, walks, gutters, and into cracks and other places where insects hide. Do not treat plants with this dosage rate.

See chart for complete insect list -- use dosages shown unless otherwise indicated. 2 tablespoons = 1 fluid ounce or 2 teaspoons. 1 cup = 8 fluid ounces or 1/2 pint. 2 tablespoons in 3 gals. water = 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals. (Note: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

WARNING: May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

Do not contaminate ornamental fishponds.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not reuse container. Destroy when empty.

Buyer's guarantee limited to label claims.

This product is highly toxic if directly exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Packed by **BONIDE CHEMICAL CO., INC., UTICA, NEW YORK 13502**

BONIDE

LAWN AND GARDEN INSECT CONTROL

w/DIAZINON 25%E

Kills certain outdoor insects which destroy lawns, roses, flowers, trees, and shrubs or annoy pets and people. It works fast and remains effective for an extended period of time.

WARNING: KEEP OUT OF REACH OF CHILDREN

**DO NOT USE IN THE HOME
NET CONTENTS 1 QUART**

DIRECTIONS

HOW TO MIX AND APPLY

Step #1: For lawns and applicable areas, measure the area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear or in advance of an expected infestation. Repeat applications as needed or as noted on the rate chart.

Step #2: Use the amount of **LAWN AND GARDEN INSECT CONTROL** and water as suggested in the rate chart.

Step #3: Spray as directed in rate chart. Use coarse drenching sprays for lawns, soil, and other surface applications. Use medium fine sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

Rate Chart for Applying **LAWN AND GARDEN INSECT CONTROL w/Diazinon 25%E**

PROBLEM AREA	PESTS	RATE AND AREA COVERED	NOTES
LAWNS Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent, Bermuda	Chinch Bug, Ground Squirrels, Weevils, Beetles, Fleas, Earwigs	2 tbsps. (1 fl. oz.) in 3 gals. of water covers 125 sq. ft. 8 fl. oz. in 24 gals. of water covers 1,000 sq. ft.	Thoroughly wet grass a few hours before applying. In the north, if chinch bugs are the only insect problem, a lower rate of application can be used: 2 tbsps. in 3 gals. of water covers 250 sq. ft.
	Millipedes	4 tbsps. (2 fl. oz.) in 3 gals. of water covers 125 sq. ft. 16 oz./24 gals. water cover 1,000 sq. ft.	
	Rhodes Grass Scale Crawlers	2 tbsps. (1 fl. oz.) per 3 gals. of water covers 100 sq. ft. 10 oz./30 gals. water covers 1,000 sq. ft.	Apply when scale crawlers first emerge.