

White thru red



BONIDE

White
Yellow
Black

BENOMYL 50% W.P.
(A Curative, Protective, Systemic action fungicide)

Black

FOR CONTROL OF LISTED DISEASES OF ROSES, FLOWERS,
ORNAMENTALS, AND TURF.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAY 11 1984

[Pictures of
diseases
controlled]



Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
4-196

EPA Est. No. 4-NY-1 EPA Reg. No. 4-196

ACTIVE INGREDIENT:

*Benomyl (Methyl 1-(butylcarbamoyl)-2-
benzimidazolecarbamate)..... 50%
INERT INGREDIENTS..... 50%

*U.S. Pat. Pending E. I. DuPont De Nemours & Co.

RECEIVED

KEEP OUT OF REACH OF CHILDREN

MAR 20 1984

CAUTION

See side panel for
additional precautionary
statements.

Registration Division (TS-767)

NET WEIGHT 4 oz. Makes up to 50 gal. or treats 2,000 sq. ft.
or of lawn.

NET WEIGHT 8 oz. Makes up to 100 gals. or treats 4,000 sq. ft.
of lawn.

BONIDE BENOMYL 50% W.P., EPA Reg. No. 4-196

SIDE PANEL



FOR INFORMATION OF THE USER
SEE THE INSTRUCTIONS
ON THE CONTAINER

1 oz. size
top right
panel

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Store in a cool, dry place, and in an area inaccessible to children or pets. Allowing product to become wet during storage may lead to certain chemical changes which will reduce the effectiveness as a fungicide.

Pesticide Disposal: Wrap container in newspaper and discard in trash.

Container Disposal: Do not reuse container. Rinse thoroughly and discard in trash.

1 oz. size
top left
panel

FOR ROOTED PLANTS

ROOTS: For control of damping-off, root rot, and other root diseases, apply to the roots of plants at the time of planting or transplanting. Use a solution of 1/2 ounce of Bonide Benomyl 50% W.P. per gallon of water.

1 oz. size
bottom
center panel

1 oz. size
bottom
right panel

FRUIT: For control of fruit rot, apply to the fruit of plants at the time of harvest. Use a solution of 1/2 ounce of Bonide Benomyl 50% W.P. per gallon of water.

FLOWERS, ORNAMENTALS, FRUIT & VEGETABLES: For control of blight, apply to the flowers, ornamentals, fruit, and vegetables of plants at the time of flowering or harvest. Use a solution of 1/2 ounce of Bonide Benomyl 50% W.P. per gallon of water.

NOTE: For control of damping-off, root rot, and other root diseases, apply to the roots of plants at the time of planting or transplanting. Use a solution of 1/2 ounce of Bonide Benomyl 50% W.P. per gallon of water.

1 oz. size
bottom
left panel

RECEIVED

MAR 20 1984

BEST DOCUMENT AVAILABLE

Registration Division (TS-707)

BONIDE BENOMYL, 50% W.P., EPA Reg. No. 4-196


4 & 8 oz. sizes only

SIDE PANEL

THUN—Dollar Spot (*Sclerotinia homeocarpa*). Large Brown
Leaf Spot (*Helminthosporium*). Apply at the rate of 2 oz. (12 level
tablespoons) per 1000 sq. ft. in sufficient water to obtain
the recommended rate, usually 2 to 5 gal. per 1000 sq. ft. Apply at
first appearance of disease and continue at 10-14 day
intervals as long as needed. When conditions are unusually
favorable for development of disease, reduce interval to 5 to 7
days. Do not graze or feed cuttings from treated areas to
livestock.

1 oz. size only

Fold under flap (glued directly to bottle)



RECEIVED

MAR 20 1984

BEST DOCUMENT AVAILABLE

Registration Division (TS-767)

300. 1189.10
17.

MAY 11 1984

Bonide Chemical Company, Inc.
2 Wurz Avenue
Yorkville, NY 13495

Attention: Bert Atsma

Gentlemen:

Subject: Benoyml 50% W.P.
EPA Registration No. 4-196
Your Letter Dated March 12, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. Add "Storage and Disposal" heading to precede all information pertinent to storage and disposal. Add the word "Storage:" to precede pertinent storage information. Revise the "Pesticide disposal statement to read "Securely wrap original container in several layers of newspapers and discard in trash."
2. Add the heading "Precautionary Statements" to precede "Hazards to Humans and Domestic Animals," the human hazard signal words, and the pertinent cautionary statements.
3. Add the heading "Environmental Hazards" to precede pertinent environmental information. Delete "Keep out of lakes, streams, and ponds." Replace with "Do not apply directly to any body of water or wetlands".

A stamped copy is enclosed for your records.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

Attachment A - The Label

CONCURRENCES

SYMBOL ▶	RD/FHB:DCR-03272:Jacobys:eg:Raven:557-2226:RD-50:4/26/84:Del.5/4/84
SURNAME ▶	
DATE ▶	