


BONIDE TOMATO- POTATO DUST



CONTROLS INSECTS AND DISEASES ON TOMATOES, POTATOES, EGG PLANT, PEPPERS, CUCUMBERS, SQUASH, PUMPKIN, MELONS, CABBAGE, BROCCOLI, CAULIFLOWER, CARROTS, TURNIPS, BEANS AND CORN AS LISTED ON SIDE PANEL.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for other cautions.

1 LB. NET WEIGHT

READ LABEL CAREFULLY INCLUDING CAUTIONS BEFORE USING

Mancozeb is a systemic fungicide which controls a wide range of fungi. It is recommended for use on a wide range of crops. With the use of Mancozeb, growers can expect to obtain protection results similar to that of commercial growers using the same chemical.

DIRECTIONS FOR USE

The amount of dust and the frequency depends upon thorough and complete application. A thin uniform film to both tops and undersides of leaves. For best results, use a sprayer with adequate reach extension, will ease labor and improve results. Small canisters may be used as a dust, by partially filling and rapidly shaking. The sides in upright position. Application when wind is very light or absent, usually early morning or evening. Do not spray directly on plants or fruit to adjacent areas. May be used as a spray by mixing two tablespoons (1 oz.) per quart or 1 pound to 1 gallon. Make dust with water. Shake or agitate sprayer to keep mixture in suspension during application.

SMALL USE TABLE		
BS. PER ACRE RATE	1 LB. WILL TREAT	4 LBS. WILL TREAT
34 lbs.	1,281 sq. ft.	5,124 sq. ft.
21 lbs.	1,039 sq. ft.	4,156 sq. ft.
14 lbs.	727 sq. ft.	2,908 sq. ft.
7 lbs.	364 sq. ft.	1,454 sq. ft.
3.5 lbs.	182 sq. ft.	727 sq. ft.
1.75 lbs.	91 sq. ft.	364 sq. ft.
0.875 lbs.	45.5 sq. ft.	182 sq. ft.

TOMATOES, POTATOES, EGG PLANT & PEPPERS: To control tomato fruit worm, fall armyworm, tomato hornworm, European corn borer, stink bug, flea beetle, squash bug, Colorado potato beetle, flea beetle, leaf hoppers, anthracnose, early and late blight, septoria leaf spot, stemphylium, grey rot, spot and late blight on potatoes, early and late blight on potatoes, anthracnose, fruit rot on eggplant and peppers.

Apply 1 lb. to 1.5 lbs. per acre on tomatoes, potatoes, or 3-6 lbs. on egg plant and peppers, depending on size of plants, every 7-10 days throughout the season up to one day before harvest. In severe anthracnose areas, apply 6-8 lbs. per acre to tomatoes. Do not use on plants under glass.

CUCUMBERS, SQUASH, PUMPKINS, MELONS, CANTALOUPE: To control pickleworm, melonworm, cucumber beetles, squash bug, flea beetle, leaf hoppers, anthracnose, downy mildew, gummy stem blight, apply 3-6 lbs. per acre on squash, melons and cantaloupes, or 4-6 lbs. on pumpkins, or 4-5 lbs. on pumpkins, when runners start to form or at first sign of disease or insects. Repeat at weekly intervals up to one day before harvest. Some varieties of melons may be sensitive to Mancozeb. Check before use.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI: To control flea beetle, harlequin bug, corn earworm, armyworm, imported cabbage worm, young cabbage loopers, alternaria leaf spot, downy mildew, apply 3-6 lbs. per acre (except cabbage 4-6 lbs. max.), brussels sprout, cauliflower and kohlrabi 5-8 lbs. max. Begin applications as soon as disease threatens or insects are present and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage loopers after heads begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CARROTS, TURNIPS: To control flea beetles, harlequin bug, leafhoppers (including six spotted leafhoppers), corn earworm, imported cabbage worm, young cabbage looper, armyworm, slug and stink bugs, cercospora leaf spot and leaf blight on carrots. Apply 3-5 lbs. per acre when disease is reported in the area or insects are present, and repeat 7-10 day intervals to day of harvest. For turnips, do not apply within seven days of harvest and remove excess residue by washing.

CORN: Sweet Florida. To control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, grasshoppers and leafhoppers, helminthosporium leaf blight. Apply 3-4 lbs. per acre when disease first threatens or insects are present and repeat at 7-10 day intervals until harvest. Do not feed forage to livestock.

For larvae in the whorl (baworm damage) and foliage feeders, apply to the entire plant. For insects attacking silks and ears, apply at two to three day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be necessary.

BEAN: To control Mexican bean beetle, leafhoppers, bean leaf beetle, cucumber beetles, western bean cutworm, flea beetles, Japanese beetles, velvet bean caterpillar, corn earworm, armyworms, stink bugs, tarnished plant bug, downy mildew, rust, apply 3-5 lbs. per acre when disease threatens or insects are present and repeat at weekly intervals until one day before harvest. - string beans, four days before harvest.

LETTUCE, ENDIVE: To control flea beetles, leafhoppers, harlequin bug, six spotted leafhopper (aster yellows vector), corn earworm, armyworms, imported cabbage worm, spittlebugs, lygus and stink bugs, downy mildew, apply 3-5 lbs. per acre when disease is reported in the area or insects are present, and repeat at 7-10 day intervals up to 14 days before harvest. - remove residues by trimming or washing.

CAUTION: Store out of reach of children. Harmful if swallowed or inhaled. Remove residue at harvest by washing or wiping. Avoid contact with and contact with skin, may cause irritation of nose, throat, eyes and skin. In case of contact, wash thoroughly with water.

Avoid contamination of foodstuffs, feed, feeding troughs and water receptacles.

Toxic to fish. Keep out of any body of water.

To protect honeybees, avoid use if possible when bees are visiting the crop. If necessary to use during such periods, warn beekeepers to move hives to a safe distance until one week after application.

Do not reuse container. Bury or dispose of in a safe place. Keep container closed when not in use and store in a cool dry place.

Buyer's guarantee limited to label claims.
Manufactured by
BONIDE CHEMICAL CO. INC. UTICA, N. Y. 13502