

BONIDE TOMATO-VEGETABLE DUST, EPA Reg. No. 4-197

VEGETABLE

USE THIS PRODUCT ONLY ON THE VEGETABLES LISTED BELOW

TOMATOES: To control leaf beetles, high quality lepidopteran and fungicide recommended for use only as indicated below. With this product the result garden can expect very productive results similar to that of commercial growers using the same chemicals.

INSTRUCTIONS FOR USE

Control of insects and diseases depends upon thorough and complete application in their vulnerable stage in both tops and underparts of leaves. Use of a good pressure sprayer with adjustable length extension will cover better and insure results. Small growers may be made as a dealer by partially washing and applying spraying the sides in regular position. Application when wind is very light or absent, usually early morning or evening, will insure best and best of results in adjacent areas. May be used as a spray by mixing with water (1 or 2 per quart, or 1 pound to 4 gallons). Make sure of that and water, strain to break up lumps, then add balance of water. Shake or agitate sprayer to keep mixture in suspension during application.

BONIDE USE TABLE		
1 LB. PER ACRE RATE	1 LB. WET TABLET	4 LB. WET TABLET
24 lbs.	1,281 sq. ft.	3,152 sq. ft.
48 lbs.	2,562 sq. ft.	6,304 sq. ft.
72 lbs.	3,843 sq. ft.	9,456 sq. ft.
96 lbs.	5,124 sq. ft.	12,608 sq. ft.
120 lbs.	6,405 sq. ft.	15,760 sq. ft.
144 lbs.	7,686 sq. ft.	18,912 sq. ft.
168 lbs.	8,967 sq. ft.	22,064 sq. ft.
192 lbs.	10,248 sq. ft.	25,216 sq. ft.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CORN: (Sweet - Florida) - To control corn earworm, corn rootworm, European corn borer, fall armyworm, flea beetles, Japanese beetle, and leafhopper, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals until harvest. Do not feed foreign to livestock. For larvae in the whorl (Chrysomelid damage) and leafhopper, apply to the entire plant. For insects attacking silks and ears, apply at two to three day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be necessary. Application of this product to the basal region of ears during the pollen shed period will seriously reduce ear population.

CORN: (Sweet - Florida) - To control corn earworm, corn rootworm, European corn borer, fall armyworm, flea beetles, Japanese beetle, and leafhopper, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals until harvest. Do not feed foreign to livestock. For larvae in the whorl (Chrysomelid damage) and leafhopper, apply to the entire plant. For insects attacking silks and ears, apply at two to three day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be necessary. Application of this product to the basal region of ears during the pollen shed period will seriously reduce ear population.

CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

CANTALOUPE: To control leaf beetles, hornworm, cabbage looper, imported cabbage worm, young cabbage looper, diamond-back moth, armyworm, cutworm, root-knot, root-rot, damping-off, black rot, bacterial wilt, and other diseases, apply 24-48 lbs. per acre when disease first develops or insects are present, and repeat at 7-10 day intervals up to three days before harvest. For small young cabbage looper after adults begin to form, apply every 5-7 days. Do not apply to cabbage within 7 days of harvest.

© 1973 & 1974 by BONIDE CHEMICAL COMPANY, INC., Washfield, N.Y. 12155

OCT 7 1982

Arnold E. Olson
Ronide Chemical Co., Inc.
2 Wurz Avenue
Yorkville, NY 13495

Dear Mr. Olson:

Subject: Ronide Tomato Vegetable Dust
EPA Registration No. 4-197
Your Application of September 9, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (78-767)

Enclosure

78-767;Ellenberger;DCR-24759;WANG-0501C;hk;Haven;479-2013;10/4/82

BEST DOCUMENT AVAILABLE



Red

White thru red-----BONIDE

4-197

White
Outline

Black
print

Yellow

TOMATO-
VEGLTABLE
Dust

CONTROLS INSECTS AND DISEASES ON TOMATOES, POTATOES,
EGG PLANT, PEPPERS, CUCUMBERS, SQUASH, PUMPKIN, MELONS,
CABBAGE, BROCCOLI, CAULIFLOWER, CARROTS, TURNIPS, BEANS
AND CORN AS LISTED ON SIDE PANEL.

100
100
100
100

BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Remove residue at harvest by washing or wiping.

STATEMENT OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. Do not induce vomiting. Do not give anything by mouth to an unconscious person.
- If inhaled - Remove victim to fresh air.
- If off skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush with plenty of water. Get medical attention immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

4
5
6
7
8
9