

Do not use on maidenhair fern, cape chestnut, or Geraldton wax flower. Excessive dosage may cause slight leaf injury to new growth on roses, phlox and petunias.

USE RECOMMENDATIONS

PLANTS	PESTS CONTROLLED	AMOUNT TO MIX WITH WATER
Ornamentals - such as roses, camellias, honeysuckle, azalias, other flowering plants, vines, privet, lilac and other herbaceous shrubs, shade trees including elm, maple, ash, locust, evergreen —such as pine, spruce, juniper, arborvitae, and various nursery plants.	Loopers, tent caterpillar, European pine shoot moth, [redacted] birch leaf miners, thrips, red-pine sawfly, lygus and other plant bugs, leafhoppers.	2 teaspoons per gal. or 1 qt. per 100 gals.
	Aphids, soft scale, scale crawlers, mealy bugs, white fly and mites.	3 teaspoons per gal. or 1½ pints per 50 gals.
Turf: bluegrass, Bermudagrass, dichondra, and other ground cover.	Sod webworms (lawn moth), cutworms, armyworms, millipeds, sowbugs (pillbugs), snails and slugs.	3 teaspoons per gal. or 1½ pints per 50 gals.

Buyer's guarantee limited to label claims and purchase price paid

Packaged by
BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502



DIRECTIONS FOR USE

Apply ZECTRAN 2E in water sprays to ornamental plants and turf. Applications may be made with hand sprayers, hose attached sprayers or power spraying equipment. Use enough spray to wet the plants and turf areas thoroughly.

KILLS: Armyworms, cutworms, sod webworms, loopers, birch leaf miners, European pine shoot moth, tent caterpillars, red-pine sawfly, thrips, leafhoppers, white fly, mealy bugs, lygus and other plant bugs, aphids, soft scales and scale crawlers, mites, millipeds, [redacted] gypsy moth, snails and slugs.

FOLIAGE-FEEDING INSECTS AND MITES: Spray infested plants thoroughly at amounts indicated in the dosage table. Re-treat as necessary.

SNAILS, SLUGS AND TURF PESTS: Water turf thoroughly before treating. Spray infested area using as indicated in the dosage table. Apply 1 gallon of spray per 130 to 180 sq. ft. of area. Soak the debris and soil under infested plants. Re-treat as necessary.

Note: Feeding stops soon after treatment, but complete kill may be delayed for two or three days.

WARNING: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. Avoid eye and skin contact. Avoid breathing spray mists. Wash well after handling. If swallowed, call a physician immediately.

Note to Physician: ZECTRAN 2E insecticide has cholinergic effects — treat symptomatically. Atropine is an antidote.

Avoid contamination of feed, foodstuffs, water supplies and fish ponds.

This product is toxic to wildlife. Do not contaminate water by cleaning equipment or disposal of waste. Bury waste in non-cropland away from water supplies. Containers should be disposed by burying with wastes.

ACCEPTED

Ornamentals - such as roses, camellias, honeysuckle, azalias, other flowering plants, vines, privet, lilac and other herbaceous shrubs, shade trees including elm, maple, ash, locust, evergreen—such as pine, spruce, juniper, arborvitae, and various nursery plants.	Loopers, tent caterpillar, European pine shoot moth, [redacted] birch leaf miners, thrips, red-pine sawfly, lygus and other plant bugs, leafhoppers.	2 teaspoons per gal. or 1 qt. per 100 gals.
	Aphids, soft scale, scale crawlers, mealy bugs, white fly and mites.	3 teaspoons per gal. or 1½ pints per 50 gals.
Turf: bluegrass, Bermudagrass, dichondra, and other ground cover	Sod webworms (lawn moth), cutworms, armyworms, millipeds, sowbugs (pillbugs), snails and slugs.	3 teaspoons per gal. or 1½ pints per 50 gals.

Buyer's guarantee limited to label claims and purchase price paid

Packaged by
BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502



DIRECTIONS FOR USE

Apply ZECTRAN 2E in water sprays to ornamental plants and turf. Applications may be made with hand sprayers, hose attached sprayers or power spraying equipment. Use enough spray to wet the plants and turf areas thoroughly.

KILLS: Armyworms, cutworms, sod webworms, loopers, birch leaf miners, European pine shoot moth, tent caterpillars, red-pine sawfly, thrips, leafhoppers, white fly, mealy bugs, lygus and other plant bugs, aphids, soft scales and scale crawlers, mites, millipeds, [redacted] gypsy moth, snails and slugs.

FOLIAGE-FEEDING INSECTS AND MITES: Spray infested plants thoroughly at amounts indicated in the dosage table. Re-treat as necessary.

SNAILS, SLUGS AND TURF PESTS: Water turf thoroughly before treating. Spray infested area using as indicated in the dosage table. Apply 1 gallon of spray per 130 to 180 sq. ft. of area. Soak the debris and soil under infested plants. Re-treat as necessary.

Note: Feeding stops soon after treatment, but complete kill may be delayed for two or three days.

WARNING: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. Avoid eye and skin