

March 5, 1973

4-201

TREATS 2,500 SQUARE FEET

A 50 Ft. by 50 Ft. Lawn Area

Clean, easy to use, lightweight granules • Just pour in spreader and go

BONIDE

BENOMYL*

LAWN FUNGICIDE GRANULES

DUPONT DE NEMOURS & CO., INC. PHILADELPHIA, PA. 19101

Provides Both Curative (Eradicant) & Residual (Protective) Action

ACTIVE INGREDIENT:

EPA Reg. No. 4-201

*Benomyl (Methyl 1-(butylcarbamoyl)-
2-benzimidazolecarbamate) 16%

INERT INGREDIENTS: 98.4%

*U.S. Pat. Pending E. I. du Pont de Nemours & Co., (Inc.), Wilmington, Delaware U.S.A.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

(See back panel cautions)

NET WEIGHT 5 LBS.

Camerman's Note
Poor Copy

BENOMYL* LAWN FUNGICIDE GRANULES is a new concept product for controlling certain lawn diseases. Now, for the first time, a fungicide has been invented which cures as well as stops certain lawn diseases. **LAWN FUNGICIDE GRANULES** made from that fungicide is a handy, convenient form for easy application by lawn spreader.

DIRECTIONS

Water lawn before application or apply after rainfall or when dew is present. Apply 2 to 5 gallons of water to each 1,000 sq. ft. If treating golf greens, always treat aprons and approaches.

For Dollar Spot (*Sclerotinia homoeocarpa*) - a summer disease - Apply contents of this bag to 2,500 sq. ft., at the rate of 2 lbs. per 1,000 square feet, at earliest appearance of disease and continue at 10-14 day intervals as long as needed.

Large Brown Patch (*Rhizoctonia solani*) - a summer disease - Apply contents of this bag to 1,250 sq. ft. at the rate of 4 lbs. per 1,000 square feet at earliest appearance of disease and continue at 10-14 day intervals as long as needed. When conditions are unusually favorable (warm & moist) for development of disease, reduce interval to 5 to 7 days.

Fusarium Patch-Pink Snow Mold (*Fusarium nivale*) - a cool season disease - Apply contents of this bag to 1,000 sq. ft. at the rate of 4 lbs. per 1,000 sq. ft. at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed.

Fusarium Blight - Frog Eye (*Fusarium roseum* and *F. tricinctum*) - a warm season disease - Apply contents of this bag to 500 sq. ft. or in severe cases, 300 sq. ft. at the rate of 10 lbs. per 1,000 sq. ft., or 16 lbs. per 1,000 in severe cases, at earliest appearance of disease and repeat 10 to 14 days later. Immediately after application, water sufficiently to thoroughly wet soil.

Striped Smut (*Ustilago striiformis*) -

Suggested Spreader Settings*

Spreader	2,500 sq. ft.	1,250 sq. ft.
Cyclone "B"	** 3-3/4	4-1/4
Jackson	8	10
Lawn Beauty	4	5
Modern	5	6
Greenfield	5	6
Patco	5	6
Scotts	4	5
Sears	5	6
Squire Applegate	5	6

*Approximate; adjustment must be made for condition of spreader, rate of walk, etc., as a prior application to smaller area indicates. Fractions estimated between whole numbers increasing numbers usually mean faster application. Close shut off on turns. Keep handle low or container vertical to ground level. Drag spreader to avoid disturbing granules after application.

**Rotary broadcast spreader for use only in absence of air movement.

NOTICE TO BUYER: Seller warrants that this product conforms to the chemical description on the label hereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, ex-

square feet, at earliest appearance of disease and continue at 10-14 day intervals as long as needed.

Large Brown Patch (Rhizoctonia solani) - a summer disease - Apply contents of this bag to 1,250 sq. ft. at the rate of 4 lbs. per 1,000 square feet at earliest appearance of disease and continue at 10-14 day intervals as long as needed. When conditions are unusually favorable (warm & moist) for development of disease, reduce interval to 5 to 7 days.

Fusarium Patch-Pink Snow Mold (Fusarium nivale) - a cool season disease - Apply contents of this bag to 1,000 sq. ft. at the rate of 4 lbs. per 1,000 sq. ft. at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed.

Fusarium Blight - Frog Eye (Fusarium roseum and F. tricinctum) - a warm season disease - Apply contents of this bag to 500 sq. ft. or in severe cases, 300 sq. ft. at the rate of 10 lbs. per 1,000 sq. ft., or 16 lbs. per 1,000 in severe cases, at earliest appearance of disease and repeat 10 to 14 days later. Immediately after application, water sufficiently to thoroughly wet soil.

Striped Smut (Ustilago striiformis) - Make a single application of the contents of this bag to 400 sq. ft. at the rate of 12 lbs. per 1,000 sq. ft. in October or early spring before grass begins growth. After application water in using 2 to 5 gallons per 1,000 sq. ft. to carry material to base of plants.

CAUTION: Avoid breathing dust, or contact with eyes or clothing. In case of contact, flush eyes with plenty of water; for eyes, get medical attention. Wash thoroughly after using. Keep away from heat, fire or sparks.

Lawn Beauty	4	5
Modern	5	5
Greenfield	5	5
Patco	5	5
Scotts	5	5
Sears	5	5
Squire Applegate	5	5

*Approximate; adjustment must be made for condition of spreader, rate of walk, etc., as a prior application to smaller area indicates. Fractions estimated between whole numbers increasing numbers usually mean faster application. Close shut off on turns. Keep handle low or container vertical to ground level. Drag spreader to avoid disturbing granules after application.

**Rotary broadcast spreader for use only in absence of air movement.

NOTICE TO BUYER: Seller warrants that this product conforms to the chemical description on the label hereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions or under abnormal use conditions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

IMPORTANT: Keep bag closed when not applying. Do not allow product to become damp or wet before use. Do not graze or feed clippings from treated areas to livestock. Do not reuse container; bury when empty.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.