

# BONIDE

TREATS 2,500 SQUARE FEET  
A 50 FOOT BY 50 FOOT LAWN AREA  
NET WEIGHT 10 LBS.

Just pour in spreader and walk.  
Clean, easy to use, dust free, dry granules.

## DIAZINON 2 $\frac{1}{2}$ G

kills bugs on lawns: weevils, lawn moths, ants, fleas, mosquitoes, bermuda and clover mites, earwigs, millipedes, fall army worms, cutworms, leafhoppers, beetles, dog ticks, crickets, chiggers, centipedes, millipedes, springtails, spiders and scorpions.

EPA Reg. No. 4-709

### ACTIVE INGREDIENT

\*0.6-diethyl 0-(2-iso-propyl-6-methyl-4-pyrimidinyl)-phosphorothioate

2.50%

### INERT INGREDIENTS

97.50%

\*Diazinon ®, a registered trademark of Ciba-Geigy Corp.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
(See back panel cautions)

# DIAZINON 2½G

**BONIDE DIAZINON 2½G** is a new and effective insecticide in dry granular form. Prevents surface feeding insect damage to lawns, and may also be used on air turf around ornamental plants, picnic areas, patios. Nothing to add! No water to mix! No spray to drift! Easily applied to lawn areas with any spreader.

## DIRECTIONS

**Begin** protection before lawn shows signs of surface insect injury. Serious damage can be prevented if the first application is made during late spring or early summer. **DIAZINON 2½G** is easily applied after mowing when grass is dry. If the infestation is very heavy, a second application will be necessary. Repeat full treatment in 7 to 10 days.

Apply at the rate of 10 lbs. per 2,500 sq. ft. or 20 lbs. per 5,000 sq. ft. Remove leaves, dead or tall grass and other debris before application. Measure area. Apply to measured area correct amount of **DIAZINON 2½G** at recommended rate with drop type spreader. Do not guess. Spread uniformly for best results. Criss cross application is recommended. Material applied in one direction, the remainder at right angles. Avoid over or under application, skips or misses.

Thoroughly wet down grass with water before application. Use at the recommended rate for each square foot.

The following suggested spreader settings are approximate. Adjustments must be made for condition of the spreader, rate of walk and humidity, etc. under conditions. Since most machine manufacturers use only whole numbers, fractions will have to be estimated between whole numbers. Increasing numbers usually equals faster rate of application. Shut off on turns, push button off when starting. Keep machine handle low or container close to ground level. Adjustments will seldom exceed 1 number.

SPREADER	MODEL	SETTING	RATE
Patco	P220 thru P224	7	10
Lawn Beauty	18 and 24	7	10
Scotts	35-8	7 ½	10
Jackson	20 and 26	7	10
Modern	360G-1	7 ½	10
*Patco Broadcast	C	5	10
*Cyclone Broadcast	B	5	10
*Brisk walking			

**CAUTION:** Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. May be absorbed through the skin. In case of contact, wash thoroughly. Do not contaminate food, foodstuffs or drinking dishes of pets. Wash after handling.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on the label. Dispose of empty container in safe place. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

BUYER'S GUARANTEE LIMITED TO LABEL CLAIMS

Manufactured by

BONIDE CHEMICAL CO., INC., UTICA, NEW YORK 13502

# DIAZINON 2½G

BONIDE DIAZINON 2½G is a new and effective insecticide in dry granular form. Prevents surface feeding insect damage to lawns, and may also be used on all turf around ornamental plants, picnic areas, patios. Nothing to add! No water to mix! No spray to drift! Easily applied to all lawn areas with any spreader.

## DIRECTIONS

Begin protection before lawn shows signs of surface insect injury. Serious damage can be prevented if the first application is made during late spring or early summer. DIAZINON 2½G is easily applied after mowing when grass is dry. If the infestation is very heavy, a second application will be necessary. Repeat full treatment in 7 to 10 days.

Apply at the rate of 10 lbs. per 2,500 sq. ft., or 20 lbs. per 5,000 sq. ft. Remove leaves, dead or tall grass and other debris before application. Measure lawn. Apply to measured area correct amount of DIAZINON 2½G at recommended rate with drop type spreader. Do not guess. Spread uniformly for best results. Criss cross application is recommended, i.e. material applied in one direction, the remainder at right angles. Avoid over or under application, skips or misses.

Thoroughly wet down grass a few hours before applying and use at the recommended rate for each square foot.

The following suggested spreader settings are approximate. Adjustments must be made for condition of the spreader, rate of walk and humidity/moisture conditions. Since most machine manufacturers use only whole numbers, fractions will have to be estimated between whole numbers. Increasing numbers usually equals faster rate of application. Close shut off on turns, push shut off open when starting. Keep machine handle low or container of machine vertical to ground level. Adjustments will seldom exceed 2 number.

SPREADER	MODEL	SETTING NO.	RATE
Patco	P220 thru P524	7½	6
Lawn Beauty	18 and 24	7½	6½
Scotts	35 - 8	7¼	6
Jackson	20 and 26	7½	6
Modern	360G - 1	7¼	6
*Patco-Rorocast	"C"	5	4