


TREES AND SHRUBS: Maple, Ash, Dogwood, Redwood, Sycamore, Birch, Elm, Hawthorn, Linden, Magnolia, Pear, Spruce, Fir, Pine, Juniper, Yew, Cypress, Cedar, Redwood, White Oak, Black Oak, Chestnut, Pecan, Walnut, Hickory, Alder, Willow	Apply to foliage, stems, and soil. Thoroughly spray underside of leaves, stems, and soil. Do not spray flowers or fruit. Do not spray on windy days. Do not spray on wet foliage. Do not spray on plants with visible damage.
WILDLIFE PESTS IN OUTSIDE AREAS:	Add 16 fl. oz. per 3 gal. of water.

See chart for complete insect list — use dosage shown unless otherwise indicated.
 2 tablespoons = 1 fluid ounce or 6 teaspoons, 1 cup = 8 fluid ounces or 1/2 pint
 2 tablespoons in 3 gal. water = 2 teaspoons in 1 gal. water or 1/2 pint in 2 1/2 gal.
 (Note: Do not use on certain ferns, poliolepta, hibiscus, papaya, pines and gardenias, and orchids since injury may occur.)
CAUTION: Harmful if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and feedstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.
 Do not contaminate ornamental fishponds.
 This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.
 Do not reuse container. Destroy when empty.
 This product is highly toxic to bees exposed to direct treatment or residues on eggs. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer's guarantee limited to label claims
 Manufactured by
BONIDE CHEMICAL CO., INC., UTICA, NEW YORK



LAWN AND GARDEN INSECT CONTROL

W/DIAZINON® 12 1/2% E

KILLS CERTAIN OUTDOOR INSECTS WHICH DESTROY LAWNS, ROSES, FLOWERS, TREES, AND SHRUBS OR ANNOY PETS AND PEOPLE. IT WORKS FAST AND REMAINS EFFECTIVE FOR AN EXTENDED PERIOD OF TIME.

EPA Reg. No. 4-211
ACTIVE INGREDIENTS:
 *O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 12.5%
 Aromatic Petroleum Derivative Solvent 67.7%
INERT INGREDIENTS: 19.8%
 *Diazinon © trademark of CIBA-GEIGY

CAUTION: KEEP OUT OF REACH OF CHILDREN
 (See side panel cautions) **1 QUART**
DO NOT USE IN THE HOME

DIRECTIONS

HOW TO MIX AND APPLY
 Step #1: For lawns and vegetables measure the area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the rate chart.
 Step #2: Use clean sprayer. Carefully measure and mix the amount of LAWN AND GARDEN INSECT CONTROL and water as suggested in the chart below.
 Step #3: Spray as directed in rate chart. Use coarse drenching sprays for lawns, soil, and other surface applications. Use medium-fine sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

RATE CHART FOR APPLYING LAWN AND GARDEN INSECT CONTROL W/DIAZINON 12 1/2% E	
PROBLEM AREA	DATE AND AREA COVERED
LAWNS: Bluestem, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Tall Fescue, Timothy, Bermuda, Zoysia, St. Augustine, Carpetgrass, Centipede, Bahama, Paspalum, Signalgrass	4 steps: (2 fl. oz.) in 3 gal. of water covers 1,000 sq. ft. 8 steps: (4 fl. oz.) in 3 gal. of water covers 1,000 sq. ft. 4 steps: (2 fl. oz.) in 3 gal. of water covers 1,000 sq. ft. in 24 gal. of water covers 1,000 sq. ft. 4 steps: (2 fl. oz.) in 3 gal. of water covers 1,000 sq. ft. in 24 gal. of water covers 1,000 sq. ft. 3 1/2 steps: in 3 gal. of water covers 1,000 sq. ft. in 24 gal. of water covers 1,000 sq. ft. 4 steps: (2 fl. oz.) per 3 gal. of water. Apply as a thorough faller spray.
ROSES AND FLOWERS: Roses, Hibiscus, Gardenias, Camellias, Azaleas, Lilacs, Forsythias, Magnolias, Spirea, Hydrangeas, Dogwoods, Apples, Peaches, Plums, Cherries, Nectarines, Apricots, Pears, Quinces, Currants, Raspberries, Blackberries, Blueberries, Raspberries, Blackberries, Blueberries, Raspberries, Blackberries, Blueberries	Apply step each creates flat coverage. For each Control, apply only step indicated in chart. Do not apply other steps. For best results, apply Control, water spray, and water spray, or other steps of Control, water spray, or other steps as indicated in chart.
PLANTS: Maple, Ash, Dogwood, Redwood, Sycamore, Birch, Elm, Hawthorn, Linden, Magnolia, Pear, Spruce, Fir, Pine, Juniper, Yew, Cypress, Cedar, Redwood, White Oak, Black Oak, Chestnut, Pecan, Walnut, Hickory, Alder, Willow	Thoroughly wet grass a few hours before spraying. In the early morning, if possible, use the early insecticide. A lower rate of application can be used: 4 steps in 3 gal. of water covers 250 sq. ft.

