


BONIDE

TREATS 4,356 SQUARE FEET

AT 5 LBS CHLORDANE PER ACRE

NET WEIGHT 10 LBS.

Just pour in spreader and walk.
Clean, easy to use, dust free, dry granules.



5% CHLORDANE GRANULES

For lawn and turf — to control white grubs, Japanese beetle larvae, earwigs, chinch bugs, sod webworms, mole crickets, chiggers, wireworms and ants.

E.P.A. Reg. No. 4-213

ACTIVE INGREDIENT:

*Technical Chlordane 5.0%

INERT INGREDIENTS: 95.0%

*equivalent to 3% Octachloro-4,
7-methanotetrahydroindane and 2% related compounds

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See rear panel cautions)

DIRECTIONS

For Turf and Lawn Insects

5% Chlordane Granules E.P.A. No. 4-213

WHITE GRUB, JAPANESE BEETLE LARVAE: Apply to soil surface in early spring or late fall at rate of 2½ to 4½ lbs. per 1,000 sq. ft. (a 20 ft. x 50 ft. area). Use higher rates for organic (black) soils. Wash insecticide into the soil to a depth of 4-6 inches with ample quantities of water.

EARWIGS: Apply to infested lawn areas in the evening during fair weather when rain is not expected at the rate of ¼ lb. per 1,000 sq. ft. Avoid wetting the treated area for at least two nights.

CHINCH BUGS (NON-RESISTANT): Water lawn thoroughly before application. Apply at the rate of 1 lb. per 1,000 sq. ft. Do not water for 2-3 days after application.

SOD WEBWORMS (LAWN MOTHS): Water lawn thoroughly before application. Apply when worms appear at the rate of 2½ lbs. per 1,000 sq. ft. Do not water after application for several days.

MOLE CRICKETS: Water lawn thoroughly before application. Apply at the rate of 2½ lbs. per 1,000 sq. ft. Water after application to carry insecticide into the soil.

CHIGGERS: Apply to lawn, ground, litter or infested vegetation at the rate of 1 lb. per 1,000 sq. ft.

WIREWORMS: Apply to lawn at rate of 2½ lbs. per 1,000 sq. ft. Wash insecticide into soil by watering same as for grubs.

ANTS: Apply liberally to ant hills at the rate of ¼ lb. per 1,000 sq. ft. and wash insecticide into the soil.

SUGGESTED SPREADER SETTINGS

These settings will give approximate starting points for spreader application. Treat a small measured area with a weighed quantity and adjust your spreader by opening or closing to obtain correct coverage. Adjustments will be necessary due to condition of spreader, rate of walk and terrain characteristics.

| SPREADER | RATE | RATE | RATE |
|--------------------|---------------------------|---------------------------|-------------------------|
| | 4½ lbs./ 1,000 sq. ft. | 2½ lbs./ 1,000 sq. ft. | 1 lb./ 1,000 sq. ft. |
| Lawn Beauty/Asgrow | 7½ | 6¼ | 5 |
| Modern/Scotts | 7¼ | 6½ | 5¼ |
| Jackson/Patco | 7¼ | 6 | 4¼ |
| Cyclone/Ortho | 5 | 4¼ | 3½ |

CAUTION:

Avoid inhalation of dust and contact with skin. In case of contact and after use, wash thoroughly with soap and water. Keep and store out of reach of children. Keep children and pets off treated areas until material is washed into soil and grass is dry. Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

BUYER'S GUARANTEE LIMITED TO LABEL CLAIMS
1972 & Mfg. by
BONIDE CHEMICAL CO., INC., UTICA, NEW YORK 13502