

# BONIDE

B O N I D E

## BENOMYL 50% W.P. LAWN & ELM TREE DISEASE CONTROL

A CURATIVE, PROTECTIVE, SYSTEMIC ACTION FUNGICIDE  
FOR PROFESSIONAL APPLICATORS TO CONTROL DOL-  
LAR SPOT AND LARGE BROWN PATCH ON LAWNS,  
AND AS AN AID IN THE CONTROL OF DUTCH ELM  
DISEASE ON ELM TREES.

EPA Reg. No. 4-215

ACTIVE INGREDIENT:

\*Benomyl (Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate 50%  
INERT INGREDIENTS: 50%

\*U.S. Pat. Pending E. I. DuPont de Nemours & Co.

IMPORTANT: Never allow BENOMYL to become wet during storage. This  
may lead to certain chemical changes which will reduce the effectiveness  
as a fungicide. Keep container closed when not in use. Do not re-use  
container; buy when empty.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

(See side panel cautions)

NET WEIGHT 8 OZ.

**Trunk injection** — Use injector tubes equipped with cups of approximately 2 fl. oz. capacity; use BENOMYL at the rate of 2 lbs. per 100 gals. water. Insert injector tubes into outer growth rings just far enough so liquid will not leak at point of entry. Space at 2 inch intervals around the trunk. Fill cups and leave in place for 24 to 48 hours; refill cups as needed. Remove injector tubes when treatment has been completed. Treatment may be made anytime during the growing season, preferably in the spring when trees reach full leaf. Repeat treatments may be made.

**CAUTION:** May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Keep away from fire or sparks. In case of contact, flush skin and eyes with plenty of water; for eyes, get medical attention.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Buyer's guarantee limited strictly to label claims

Packed by

BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502

**TURF** — Dollar Spot (*Sclerotinia homocarpa*), Large Brown Patch (*Rhizoctonia solani*): Apply at the rate of 2 oz. (12 level tablespoons per 1,000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1,000 sq. ft.) Apply at earliest appearance of disease and continue at 10-14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce intervals to 5 to 7 days.

**ELM TREES** — Dutch Elm Disease: Foliar Spray — Use BENOMYL at the rate of 8 lbs. per 100 gals. water, and apply 2 to 3 gals. of spray per mature tree using a concentrate sprayer. A surfactant may be added to the spray mixture to improve wetting of foliage. Apply in the spring when trees reach full leaf, which is the period when elm bark beetles usually begin feeding.

### DIRECTIONS

ACCEPTED

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
EPA UNDER NO. 4-215