

B O N I D E

INSECT BAIT GRANULES
W/SEVIN

CONTROLS CUTWORMS, ARMYWORMS, CRICKETS, DARKLING
GROUND BEETLES, GRASSHOPPERS AND SOWBUGS

EPA Est. 4-NY-1
EPA Reg. No. 4-232

ACTIVE INGREDIENT:

*Carbaryl (1-naphthyl methylcarbamate)..... 5%

INERT INGREDIENTS:.....95%

*SEVIN is the registered T.M. of the
Union Carbide Corporation

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if inhaled or swallowed. Avoid breathing of dust and contact with skin. Do not take internally. Wear regular long sleeved clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, streams and ponds. Keep pets off treated areas.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

Buyer's guarantee limited to label claims

25 LBS. NET WEIGHT



BONIDE INSECT BAIT W/SEVIN provides prompt and lasting control of cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs. These insect pests often work close to the ground, sometimes under heavy crop vegetative cover. Sprays and dusts may not reach them, or provide lasting effect. BONIDE INSECT BAIT drops to the ground through heavy cover, as well as on bare soil, in exact position to attract and destroy insects before they can damage your crops. They fit perfectly into many integrated insect control programs for destructive insects. The baits destroy damaging insects without effect on many beneficial insects; can be used without residue problems on a great variety of crops, and you have no drift problems at all. Just follow label directions.

DIRECTIONS FOR USE


Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20 to 40 pounds per acre of bait will control cutworms*, armyworms, crickets, darkling ground beetles, grasshoppers and sowbugs.

Crop	Amount	Days Between Final Application and Harvest
cucumbers, melons, peanuts	20 lbs./acre	None
alfalfa, peas	30 lbs./acre	None
cotton	30 lbs./acre	None
beans, carrots, corn, eggplants, okra, peppers, potatoes, tomatoes, sweet corn	40 lbs./acre	None
asparagus, strawberries	40 lbs./acre	1 day
broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, beets (roots), horseradish, turnips, radishes, rutabaga, turnips (roots)	40 lbs./acre	3 days
sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard, turnip (tops)	40 lbs./acre	14 days

*For cutworms that work at night, spread bait in the afternoon, banded along rows or broadcast.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residue when SEVIN is used in accordance with label directions. If this product is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn cobs and cotton bolls may be grazed or harvested for use as food for dairy and meat animals.



MAY 7, 1973

1 LBS



SUPER Blue Dragon



GARDEN DUST

5% SEVIN

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects.

ACTIVE INGREDIENT	BY WEIGHT
Carbaryl (1 naphthyl N methylcarbamate)	5%
INERT INGREDIENTS	95%
TOTAL	100%

CAUTION:

EPA REG. NO. 16-15

Manufactured By

DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24061



SUPER BLUE DRAGON

NEAREST NEIGHBOR
CALLING NUMBER
FEDERAL AGENCY

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

SEVIN

BLUE DRAGON



SUPER