


**TREATS 2,500 SQUARE FEET
A 50 FOOT BY 50 FOOT LAWN AREA
NET WEIGHT 3 LBS. 8 OZS.**

**Just Pour in Spreader and Walk
Clean, Easy to Use, Dry Granules**



**LAWN
WEED KILLER
GRANULES**

Comments

WEED KILLER

GRANULES

KILLS

Dandelion, Plantain, buckhorn, thistle, sow thistle, ground ivy, ragweed, burdock, nettle, pennywort, pigweed, healall, sheep sorrell, yarrow, wild onion, purslane, chicory, gypsum weed, black medic and white clover in established lawns.

E.P.A. EST. 4-NY-1 E.P.A. Reg. No. 4-243

ACTIVE INGREDIENTS:

*Triethanolamine 2,4-Dichlorophenoxyacetate.....	4.69%
**Triethanolamine Silvex (2-(2,4,5-trichlorophenoxy)propionic Acid).....	1.26%

INERT INGREDIENTS: 94.05%

*equivalent to 2,4-D Acid 2.79%

**equivalent to 2,4,5-TP acid .74%

CAUTION: KEEP OUT OF REACH OF CHILDREN
(Note back panel Cautions)

LAWN WEED KILLER

GRANULES

Lawn Weed Killer can be applied any time spring, summer or fall for general weed control. Application during warm periods 60-85° when weeds are actively growing gives best results. During cool or cold periods action is slow and retreatment may be necessary. Do not apply to a new lawn until after second regular mowing.

DIRECTIONS

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a **light** shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for $\frac{1}{4}$
inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded two weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Storage in cool dry area but not near seeds, fertilizers, insecticides or fungicides prolongs effectiveness of ingredients. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds or on Bermuda and St. Augustine grasses. May cause injury to bentgrass, Dichondra, or carpetgrass lawns. **Do not contaminate water used for irrigation or domestic purposes.**

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for $\frac{1}{4}$
inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded two weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Storage in cool dry area but not near seeds, fertilizers, insecticides or fungicides prolongs effectiveness of ingredients. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds or on Bermuda and St. Augustine grasses. May cause injury to bentgrass, Dichondra, or carpetgrass lawns. Do not contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy it by burying in a safe place away from water supplies. Do not burn.

Starting Point suggested spreader settings:

Make & Model	Setting	Make & Model	Setting
Scott	8	Montgomery Ward 89-1081	$8\frac{3}{4}$
Jackson 18-26 18"	7	Salem Micro Set 22"	7
Sears, Roebuck Cross Country 19"	7	Cyclone B 5' Swath (Still Air Only)	6
Lawn Beauty	7		

**TREATS 2,500 SQUARE FEET
A 50 FOOT BY 50 FOOT LAWN AREA
NET WEIGHT 3 LBS. 8 OZS.**

**Just Pour in Spreader and Walk
Clean, Easy to Use, Dry Granules**



LAWN

WEED KILLER

GRANULES

registered under
EPA Reg. No. 4243

Lawns

LAWN

WEED KILLER

GRANULES

KILLS

Dandelion, Plantain, buckhorn, thistle, sow thistle, ground ivy, ragweed, burdock, nettle, pennywort, pigweed, healall, sheep sorrell, yarrow, wild onion, purslane, chicory, gypsum weed, black medic and white clover in established lawns.

E.P.A. EST. 4-NY-1 E.P.A. Reg. No. 4-243

ACTIVE INGREDIENTS:

- *Triethanolamine 2,4-Dichlorophenoxyacetate..... 4.69%
- **Triethanolamine Silvex (2-(2,4,5-trichlorophenoxy)propionic Acid)..... 1.26%

INERT INGREDIENTS: 94.05%

*equivalent to 2,4-D Acid 2.79%
**equivalent to 2,4,5-TP acid .74%

CAUTION: KEEP OUT OF REACH OF CHILDREN
(Note back panel Cautions)

LAWN WEED KILLER

GRANULES

Lawn Weed Killer can be applied any time spring, summer or fall for general weed control. Application during warm periods 60-85° when weeds are actively growing gives best results. During cool or cold periods action is slow and retreatment may be necessary. Do not apply to a new lawn until after second regular mowing.

DIRECTIONS

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for $\frac{1}{4}$
inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded two weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Storage in cool dry area but not near seeds, fertilizers, insecticides or fungicides prolongs effectiveness of ingredients. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds or on Bermuda and St. Augustine grasses. May cause injury to bentgrass, Dichondra, or carpetgrass lawns. Do not contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy it by burying in a safe place away from water supplies.

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.

2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.

3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for 1/4 inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.

4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.

5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded two weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

CAUTION

ACCEPTED WITH COMMENTS

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Storage in cool dry area but not near seeds, fertilizers, insecticides or fungicides prolongs effectiveness of ingredients. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds or on Bermuda and St. Augustine grasses. May cause injury to bentgrass, Dichondra, or carpetgrass lawns. Do not contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy it by burying in a safe place away from water supplies. Do not burn.

Starting Point suggested spreader settings:

Make & Model	Setting	Make & Model	Setting
Scott	8	Montgomery Ward 89-1081	8 ³ / ₄
Jackson 18-26 18"	7	Salem Micro Set 22"	7
Sears, Roebuck Cross Country 19"	7	Cyclone B 5' Swath (Still Air Only)	6
Lawn Beauty	7		

7

**TREATS 2,500 SQUARE FEET
A 50 FOOT BY 50 FOOT LAWN AREA
NET WEIGHT 3 LBS. 8 OZS.**

**Just Pour in Spreader and Walk
Clean, Easy to Use, Dry Granules**

4-243

LAWN WEED KILLER

GRANULES

KILLS

Dandelion Plantain buckhorn thistle sow thistle ground ivy ragweed
burdock nettle pennywort pigweed healall sheep sorrell yarrow wild
onion parsnip cherry oxysum weed black medic and white clover
in established lawn

© 1974 by The Scotts Company, Kenilworth, N. J. 07033

E. 4-243

LAWN WEED KILLER

GRANULES

KILLS

Dandelion, Plantain, buckhorn, thistle, sow thistle, ground ivy, ragweed, burdock, nettle, pennywort, pigweed, healall, sheep sorrell, yarrow, wild onion, purslane, chicory, gypsum weed, black medic and white clover in established lawns.

E.P.A. EST. 4-NY-1 E P.A. Reg. No. 4-243

ACTIVE INGREDIENTS:

*Triethanolamine 2,4-Dichlorophenoxyacetate	4.69%
**Triethanolamine Silvex (2-(2,4,5-trichlorophenoxy)propionic Acid)	1.26%
INERT INGREDIENTS:	94.05%

*equivalent to 2,4-D Acid 2.79%
**equivalent to 2,4,5-TP acid .74%

CAUTION: KEEP OUT OF REACH OF CHILDREN
(Note back panel Cautions)

LAWN WEED KILLER

GRANULES

Lawn Weed Killer can be applied any time spring, summer or fall for general weed control. Application during warm periods 60-85° when weeds are actively growing gives best results. During cool or cold periods action is slow and retreatment may be necessary. Do not apply to a new lawn until after second regular mowing.

DIRECTIONS

1. Mow lawn 2-4 days prior to application and delay mowing for three days after application.
2. Choose a calm period (to avoid wind drift of material to valuable plantings) when rain is not expected for 24 hours. Early morning when lawn is moist with dew or after a light shower or hose watering is the best time to apply.
3. Measure area to be treated and apply contents of this bag to area listed on front of bag. Spreader settings for 1/4 inch wide openings will work on most spreaders but due to wear and different walking rates, adjust your spreader to avoid gross under or over dose. Other suggested spreader settings on list below. Avoid overlapping and double coverage in backing up or turning, close spreader at those times and before filling spreader. In case of accidental heavy overdose, watering the overdosed area will reduce or prevent damage.
4. Do not water the lawn after application. Twisting and loss of color of weeds within 7 days indicates effectiveness.
5. It is not essential that dead or dying weeds be raked out. Bare spots may be reseeded two weeks after application. 2 to 3 days prior to reseeding, heavily water the lawn.

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact and after use wash or flush with water.

Avoid contamination of feed, foodstuffs, fish ponds, water supplies and streams. Storage in cool dry area but not near seeds, fertilizers, insecticides or fungicides prolongs effectiveness of ingredients. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not use on flowers, vegetables or ornamental beds or on Bermuda and St. Augustine grasses. May cause injury to bentgrass, Dichondra, or carpetgrass lawns. Do not contaminate water used for irrigation or domestic purposes.

Do not reuse container. Destroy it by burying in a safe place away from water supplies. Do not burn.

Starting Point suggested spreader settings

Make & Model	Setting	Make & Model	Setting
Scott	8	Montgomery Ward 89-1081	8 1/2
Jackson 18 26 18'	7	Salem Micro Set 22"	7
Ward 18 26 18'	7	Ward 18 26 18'	6
Ward 18 26 18'	7	Ward 18 26 18'	6