

4-251


6/1/2004

Page 1 of 3



DIPEL 150 DUST

(*Bacillus thuringiensis*)



FOR VEGETABLES

Optional Marketing Statements:

Ready-to-Use

A Biological Insecticide Controls

Cabbage Looper, Imported Cabbageworm, Grape Leaf Folder and Hornworm on Crops Listed on Side Panel.

Easy to Use Duster Method.

ACTIVE INGREDIENT:

Bacillus thuringiensis, subspecies *kurstaki*, strain ABTS-351,
fermentation solids and solubles 0.47%
(16,000 International Units of Potency per mg)

OTHER INGREDIENTS: 99.53%

TOTAL 100.00%

*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side/back/other panel(s) for additional Precautionary Statements

NET WEIGHT: xx LB (xx g)

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Avoid breathing dust.

Environmental Hazards - Keep out of lakes, streams or ponds. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For residential use in vegetable gardens and on fruits, shade trees and ornamentals.

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Disposal: *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Dipel 150 Dust is a biological insecticide intended for use against chewing larvae (caterpillars) of certain lepidopterous species. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

CROP	INSECT	APPLICATION RATE
Broccoli, Cabbage, Cauliflower, Collards, Celery, Melons, Potatoes, Spinach and Cucumbers	Diamondback Moth Hornworms Cabbage Looper Melonworms Webworm Green Cloverworm	4 to 8 oz per 435 sq. ft.
Broccoli, Cabbage, Cauliflower, Collards, and Turnip Greens	Loopers Imported Cabbageworm Hornworms	2½ to 4 oz per 435 sq. ft.
Grapes, Small fruits, Cane and Bush Berries	Loopers Omnivorous Leaf Rollers Grape Leaf Folder Grape Leaf Skeletonizer	4 to 8 oz per 435 sq. ft.
Tomatoes, Peppers, Eggplant	Loopers Hornworms Tomato Fruitworm* Variegated Cutworm	2 to 4 oz per 435 sq. ft.

Lawns	Sod Webworm	Apply 4 to 8 oz per 100 sq. ft.
Flowers and Ornamentals	Loopers Tobacco Budworm Omnivorous Looper Omnivorous Leaf Roller Diamondback Moth Armyworms** Ello Moth (Hornworm) Lo Moth Oleander Moth	Apply 2 to 4 oz per 435 sq. ft.

*Apply weekly in a preventative program.

**Dipel Dust may be used to suppress small worms (1st & 2nd instar) when populations are light and full ground coverage can be attained.

Buyer's Guarantee Limited to Label Claims

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents) call National Pesticide Information Center at 1-800-858-7378.

Manufactured by:

Bonide Products, Inc.

Oriskany, NY 13424

