

BONIDE

GARDEN & ORNAMENTAL FUNGICIDE 75% WP CONTAINS: CHLOROTHALONIL

**CONTROLS LEAF SPOTS, CERTAIN MILDEWS,
CERTAIN BLIGHTS, ANTHRACNOSE,
AND OTHER LISTED DISEASES
ON VEGETABLES AND ORNAMENTALS**

EPA REG No 4-289 EPA EST No 4 NY-1

ACTIVE INGREDIENTS
Chlorothalonil (tetrachloro-sophtalonyl) 75%
INERT INGREDIENTS 25%
TOTAL 100%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

**KEEP OUT OF REACH OF CHILDREN
DANGER**

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

NET WEIGHT 1 LB.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS — DANGER

Corrosive: causes severe eye damage.
Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.
Avoid contact with skin or clothing. Wash exposed areas of skin with soap and arm water after handling or using.
Do not take internally.
Avoid breathing dust or spray mist.

Note: Bonide Garden & Ornamental Fungicide 75% WP may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having a severe allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS —

This product is toxic to fish. Keep out of lakes, streams or ponds.
Do not apply where runoff is likely to occur.
Do not apply when weather conditions favor drift from areas treated.
Apply only to areas specified on label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

GENERAL INFORMATION

Bonide Garden & Ornamental Fungicide 75% WP is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases. Bonide Garden & Ornamental Fungicide 75% WP can be used effectively in dilute or concentrate sprays. Thorough uniform coverage is essential for disease control.

Do not combine Bonide Garden & Ornamental Fungicide 75% WP with spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination phytotoxic and non-injurious under your conditions of use.

The reduced amount of Bonide Garden & Ornamental Fungicide 75% WP should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of Bonide Garden & Ornamental Fungicide 75% WP in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operation. Do not use on greenhouse grow crops.

Apple Blight — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when blossoms are in full bloom. Do not apply within 7 days of harvest. Do not spray treated areas or feed treated plant parts.

Apple Scab — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when blossoms are in full bloom. Do not apply within 7 days of harvest. Do not spray treated areas or feed treated plant parts.

Apple Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when blossoms are in full bloom. Do not apply within 7 days of harvest. Do not spray treated areas or feed treated plant parts.

Bacterial Blight — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Bacterial Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Bacterial Wilt — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Rot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Black Spot — Use 1.1 to 2.2 lbs. per acre (1.2 to 2.2 gal. of 75% WP) in sufficient water to obtain adequate coverage. Start applications when disease treatment is required. Repeat applications at 7 to 10 day intervals or as necessary to maintain control.

Buyer's guarantee limited to label claims and refund of purchase price
MFG. by BONIDE CHEMICAL CO., INC., YORKVILLE, NY 13495

62