

BONIDE

TURF, GARDEN & ORNAMENTAL FUNGICIDE 50% WP

CONTAINS: CHLOROTHALONIL

CONTROLS LEAF SPOTS, CERTAIN MILDEWS, CERTAIN BLIGHTS, ANTHRACNOSE, AND OTHER LISTED DISEASES ON TURF, VEGETABLES, FRUITS, AND ORNAMENTALS AS LISTED ON THIS LABEL.

EPA REG. NO. 4-290 EPA EST. NO. 4-NV-1

ACTIVE INGREDIENTS:
*Chlorothalonil (tetrachlorophthalonitrile) 50%
INERT INGREDIENTS 50%
TOTAL 100%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

KEEP OUT OF REACH OF CHILDREN
DANGER
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 1 LB.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - DANGER

Corrosive. Causes severe eye damage.
Do not get in eyes, wear goggles or eye shield when handling this product.
Avoid contact with skin or clothing.
Wash thoroughly with soap and water.
Do not eat, drink, or smoke while using this product.

STATEMENT OF PRACTICAL TREATMENT

If eyes are contaminated - Flush with plenty of water immediately for 15 minutes.
If on skin - Wash exposed areas of skin with soap and warm water after handling or using.

Note: Bonide Turf, Garden & Ornamental Fungicide 50% WP may produce temporary allergic effects of a rash, fever, or redness of the eyes, mild bronchial irritation and redness of skin on exposed areas. Persons having allergic reactions should contact a physician. Antihistamines are useful in treatment with antihistamines or steroid cream on the skin.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds.
Do not apply where it will drift onto crops.
Do not apply where it will drift onto areas treated.
Apply only to areas specified on label.

DIRECTIONS FOR USE

If a violation of Federal laws or this product is a danger inconsistent with this labeling.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Turf - Apply 2.14 to 3.21 lbs per 1000 sq ft of turf area. Use 2.14 lbs per 1000 sq ft for preventive control and 3.21 lbs per 1000 sq ft for curative control. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Golf Courses - Apply 2.14 to 3.21 lbs per 1000 sq ft of green area. Use 2.14 lbs per 1000 sq ft for preventive control and 3.21 lbs per 1000 sq ft for curative control. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Vegetables - Apply 2.14 to 3.21 lbs per 1000 sq ft of vegetable area. Use 2.14 lbs per 1000 sq ft for preventive control and 3.21 lbs per 1000 sq ft for curative control. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Fruits - Apply 2.14 to 3.21 lbs per 1000 sq ft of fruit area. Use 2.14 lbs per 1000 sq ft for preventive control and 3.21 lbs per 1000 sq ft for curative control. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Apply 2.14 to 3.21 lbs per 1000 sq ft of ornamental area. Use 2.14 lbs per 1000 sq ft for preventive control and 3.21 lbs per 1000 sq ft for curative control. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Disease	Rate of Application	Rate of Control	Rate of Cure
Downy Mildew	2.14	3-6	9-12
Leaf Spot	2.14	3-6	6-12
Gray Blight	2.14	3-6	6-12
Anthracnose	2.14	3-6	6-12
Large Blight	2.14	3-6	6-12
Red Blight	2.14	3-6	6-12
Downy Mildew	2.14	3-6	9-12
Anthracnose	2.14	3-6	9-12
Leaf Spot	2.14	3-6	9-12

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Ornamentals - Bonide Turf, Garden & Ornamental Fungicide 50% WP is recommended for the control of a number of important ornamental diseases and is also effective against many common lawn diseases. Apply at 7 to 10 day intervals. Do not apply to areas where it will drift onto crops.

Buyer's guarantee limited to label claims and refund of purchase price.
MFG. by BONIDE CHEMICAL CO., INC., YORKVILLE, NY 13495