

BONIDE

TREATS 5000 SQUARE FEET — Preventative
(A 100 FOOT BY 50 FOOT LAWN AREA)

TREATS 2500 SQUARE FEET — Curative
(A 50 FOOT BY 50 FOOT LAWN AREA)

NET WEIGHT 20 LBS.

TURF FUNGICIDE 5G

CONTAINS CHLOROTHALONIL

Controls leaf spots, copper spot, dollar spot,
certain mildews, certain blights, and other listed
diseases on turf

EPA EST. NO. 4-NY-1 EPA REG. NO. 4-291

ACTIVE INGREDIENT:

*Chlorothalonil (tetrachloroisophthalonitrile)

INERT INGREDIENTS

TOTAL

5
95
100

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS AND DIRECTIONS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS **DANGER**

Do not get this product on your skin or clothes. Wash immediately with plenty of water.

Avoid contact with eyes.

Do not take internally.

Avoid breathing dust or spray.

STATEMENT OF PRACTICAL TREATMENT

Eyes are contaminated — Flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

On skin — Wash exposed areas of skin with plenty of water after handling or using.

Note: Bonide Turf Fungicide may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of areas frequented by fish.

Do not apply where runoff is likely to occur.

Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

DISPOSAL

Disposal: Do not reuse empty containers. Do not dump into water.

ORNAMENTAL TURF

This product is applied with a backpack sprayer or pump sprayer. Apply to the turf area to be treated. Do not apply to other areas.

Apply to the turf area to be treated. Do not apply to other areas. Do not apply to other areas.

Apply to the turf area to be treated. Do not apply to other areas.

Disease	When to apply	Interval of Application	Preventive* before disease occurs)	Curative** (when disease first appears)
Leaf spot	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-4 bs
Leaf spot	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-4 bs
Leaf spot	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-4 bs
Leaf spot	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-4 bs
Helminthosporium leaf spot and melting out	Early spring and continue as long as wet weather prevails	7-10 day	1-4 bs	1-8 bs
Large brown patch	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-8 bs
Red thread	Early spring to early summer and continue to early fall	7-10 day	1-4 bs	1-8 bs
Leaf rust of blue grass	Early spring to early summer and continue to early fall	7-10 day	1-6 bs	6-8 bs
Dothidea Alternaria leaf spot (California only)	Early spring to early summer and continue to early fall	7-14 day	1-6 bs	6-8 bs

*Recommended rates for preventive use are based on 100% disease control when disease conditions are light to moderate. Higher indicated rates will be required for severe conditions.

**Rates for use on a curative basis when disease conditions are moderate to severe should be 10-20 lbs per acre under severe conditions.

Buyer's guarantee limited to label claims and refund of purchase price.

Manufactured by

BONIDE CHEMICAL CO., INC. • YORKVILLE, NY 13495