

183
11

BONIDE

TREATS 5000 SQUARE FEET — Preventative
(A 100 FOOT BY 50 FOOT LAWN AREA)

TREATS 2500 SQUARE FEET — Curative
(A 50 FOOT BY 50 FOOT LAWN AREA)

NET WEIGHT 20 LBS.

TURF FUNGICIDE 5G

CONTAINS CHLOROTHALONIL

Controls leaf spots, copper spot, dollar spot,
certain mildews, certain blights, and other listed
diseases on turf

EPA EST. NO. 4-NY-1 EPA REG. NO. 4-29 JUL 25 1985

ACCEPTED
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, this pesticide
registered under
EPA Reg. No. 4-291

ACTIVE INGREDIENT:			
*Chlorothalonil (tetrachloroisophthalonitrile)			5%
INERT INGREDIENTS			95%
		TOTAL	100%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Statement of Practical Treatment

Corrosive, causes severe eye damage. Skin sensitizer. Do not get in eyes, wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately. Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using. Do not take internally. Avoid breathing dust or spray mist.

First Aid: Note to physician - Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact you State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Do not reuse empty container. Wrap container and put in trash collection.

GENERAL DIRECTIONS

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

GENERAL DIRECTIONS (Cont'd)

ORNAMENTAL TURF

This product is applied with a spreader. Spreader must be properly adjusted, calibrated, and checked for accuracy before applying. Carefully measure area to be treated to determine amount of product to apply.

Do not mow or water after treatment. A 24 hour waiting period following treatment is preferred. Bonide Turf Fungicide 5G should always be used in conjunction with good turf management practices.

Do not graze treated areas. Do not feed clippings to livestock.

Disease	When to apply	Interval of Application	Rate Pounds per 1,000 sq. ft.	
			Preventive* (before disease occurs)	Curative** (when disease first appears)
Copper Spot	Early spring to early summer and continue to early fall	7-10 day	4-6 lbs.	6-8 lbs.
Curvularia leaf spot	Early spring and continue as long as wet weather prevails	7-10 day	2-4 lbs.	4-8 lbs.
Dollar spot	Early spring to early summer and continue to early fall	7-10 day	2-4 lbs.	4-8 lbs.
Grey leaf spot	Early spring to early summer and continue to early fall	7-10 day	2-4 lbs.	4-8 lbs.
Helminthosporium leaf spot and melting out	Early spring and continue as long as wet weather prevails	7-10 day	2-4 lbs.	4-8 lbs.
Large brown patch	Early spring to early summer and continue to early fall	7-10 day	2-4 lbs.	4-8 lbs.
Red thread	Early spring to early summer and continue to early fall	7-10 day	2-6 lbs.	6-8 lbs.
Stem rust of blue grass	Early spring to early summer and continue to early fall	7-10 day	4-6 lbs.	6-8 lbs.
Dichondra Ailernaria leaf spot (California only)	Early spring to early summer and continue to early fall	7-14 day	4-6 lbs.	6-8 lbs.

*Recommended rates for preventing disease establishment, use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe

**Rates for use on a 7 day schedule when disease is present. Higher indicated rate should be applied under severe conditions

Buyer's guarantee limited to label claims and refund of purchase price.

Manufactured by



BONIDE CHEMICAL CO., INC. • YORKVILLE, NY 13495