

BONIDE

FRONT PANEL

173

PM 04

4-302

11/22/88

ACCEPTED
NOV 22 1988

FLEA BEATER

4-302


BEST IMAGE AVAILABLE

KILLS FLEAS, TICKS, ANTS, CRICKETS, SPIDERS, EARWIGS AND OTHERS LISTED


Bonide FLEA BEATER provides residual pest control in and around homes, garages, trailers, campers, schools, hospitals, recreational areas . . . Easy to apply right from the container. Treats 1000 sq. ft.



Flea



Tick



Field
Cricket



Firebrat

EPA Est. 4-NY-1

EPA Reg. No. 4-302

ACTIVE INGREDIENT

Bendiocarb. (2,2-dimethyl-1,3-benzodioxol-4-yl methylcarbamate)

INERT INGREDIENTS

Formulated in U.S. Patent No. 3,726,338

1%

99%

100%

KEEP OUT OF REACH OF CHILDREN

12 pt. type

CAUTION

18 pt. type

See side panel for additional
precautionary statements and first aid.

NET WEIGHT 2.75 lbs.

282

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Do not breathe dust. Do not allow to contact skin, eyes, or clothing.

Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swallowed— Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled —Keep patient strictly at rest and transfer to hospital immediately to get medical attention.

If on skin —Wash with soap and warm water. **Seek medical attention if irritation persists**

If in eyes —Rinse thoroughly with clean water.

NOTE TO PHYSICIAN: Contains anticholinesterase carbamate insecticide. In cases of overexposure, atropine sulfate—2 mg. (equivalent to 1.30 grain adult dose) should be given and repeated as necessary until fully atropinized.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water ~~by cleaning of equipment or disposal of waste~~ when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Collect, cover or incorporate product spilled on soil surface.

PRECAUTIONS

- Do not use on or contaminate fruit, vegetables or other food or feed crops.
- Do not apply to humans, animals, clothing or bedding.
- Do not contaminate feed or food products or food preparation surfaces, dishes, kitchen utensils and food containers.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BONIDE FLEA BEATER is for residual pest control in and on buildings and structures and their immediate surroundings. Permitted areas of use include, but are not limited to, houses, garages, trailers, campers, storage sheds, apartment buildings, industrial buildings, schools, nursing homes, hospitals, hotels, and non-food areas of stores, warehouses, and restaurants.

Application: Apply FLEA BEATER thoroughly and uniformly to the area to be treated. The amount applied will vary with the site, but should usually be in the range of 1/4 to 3/4 oz. per sq. yard or 2 1/2 to 4 1/2 lbs. per 1,000 sq. ft.

INDOOR USE — BONIDE FLEA BEATER is intended for application to hiding and runway areas and those places where pests are found.

Cockroaches, Ants, Crickets, Silverfish and Firebrats: Thoroughly dust all areas where these pests crawl and hide, especially cracks and crevices around sinks, stoves, refrigerators and cabinets; behind baseboards; around doors and windows and in attics, basements, and crawl spaces. For ants, apply to ant trails and around doors and windows and other places where ants enter premises.

Carpenter Ants: Dust around doors and windows and other places where these pests enter premises. When possible, locate and dust nests and surrounding areas.

Brown Dog Ticks and Fleas in Buildings: Dust infested areas such as pet beds and resting quarters, nearby cracks and crevices, under the edges of rugs and floor coverings, between and under cushions of upholstered furniture and other places where these pests may hide. Old bedding of pets should be removed after treatment of pet area and replaced with clean fresh bedding. Additionally, ensure that pets are treated with an EPA approved product registered for pet use. **DO NOT TREAT ANIMALS OR THEIR BEDDING WITH THIS PRODUCT.**

303

Instructions for Killing Fleas in Carpets with BONIDE FLEA BEATER:
 Sprinkle lightly and evenly onto carpet. Drive Flea Beater into the carpet by brushing with a broom. Wait 2 - 4 hours then vacuum carpet. Wait 6 - 8 hours for heavy infestations. Remove bag from vacuum, wrap in newspaper and discard in trash outside the home. Carpets are chemically treated to resist stains. This means they also resist liquid flea sprays. FLEA BEATER gets into the nap of the carpet where fleas hide.

Spiders: Dust along and behind baseboards, window and door frames, corners, pipes, storage localities, attics, basements, crawl spaces and other areas where spiders may hide.

Wasps: It is best to treat wasp nests at night when insects are less active and have returned to the nest. Do not wear loose fitting clothing—take necessary precautions to avoid stings. For best results, carefully dust near nest and check one or two days after treatment to ensure complete kill, and retreat if necessary. If possible, remove and destroy nest to prevent hatching of new insects. For ground nests, carefully dust entrance and surrounding areas. Retreat in one or two days if wasp activity is still present. Do not use this product to treat overhead nests.

Sowbugs, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Apply around doors, windows and other places where these pests may enter. Treat baseboards, storage areas, and cracks and crevices, especially in dark, musty areas where these pests may hide.

OUTDOOR USE—Use BONIDE FLEA BEATER as a residual treatment for control of ants, brown dog ticks, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders and wasps. Make a residual treatment around windows and doors, porches, screens, eaves, patios, garages, under stairways and in crawl spaces and other areas where pests hide.

To help prevent invasion of buildings and structures by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building or structure. Also treat the building foundation and crawl spaces where pests are active and may find entrance. Apply thoroughly and uniformly to the band area. Pay particular attention to ant hills and wasp nests in the band area. Wasp nests in trees and shrubs may be treated as noted above.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry area inaccessible to children and pets.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Pesticide Disposal: Securely wrap original container in several layers of paper and discard in trash.

12 pt.
typ

Warranty Limited to Label Claims and Refund of Purchase Price.

BONIDE CHEMICAL CO., INC. • YORKVILLE, NY 13495