

4-303
2/21/16
2/10/4

RESTRICTED USE PESTICIDE
For retail sale to, and use only by,
Certified Applicators or persons under
their direct supervision and only for
those uses covered by the Certified
Applicator's certification.

Not for Sale or Use in California,
Texas, Utah or Puerto Rico.
BONIDE

ACCEPTED
AUG 8 1984
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 4-303

ZINC PHOSPHIDE BAIT

A RODENTICIDE FOR USE IN ORCHARDS,
FOREST NURSERIES, CHRISTMAS TREE NURSERIES,
ORAGE CROPS, NURSERIES AND LANDSCAPE
PLANTINGS. ALSO FOR USE AROUND LAWNS,
ORNAMENTALS, BUILDINGS, GOLF COURSES
AND PARKS

EPA Est. No. 4-NY-1

EPA Reg. No. 4-303

ACTIVE INGREDIENT:
Zinc Phosphide..... 1.00%
INERT INGREDIENTS: 99.00%


BEST AVAILABLE COPY

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements

NET WEIGHT 50 LBS.



Reg # 4-303
1414
2001

BONIDE ZINC PHOSPHIDE BAIT, EPA Reg. No. 4-303

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION

Keep away from humans, domestic animals and pets. May be fatal or harmful if swallowed or inhaled. Avoid skin contact. Do not breathe dust. Avoid contamination of feed or foodstuffs. Avoid contact with acids.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Avoid use of all oils. Have patient lie down and keep warm. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If on skin or in eyes: Flush with plenty of water.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and fish. Birds and other wildlife feeding on treated bait may be killed. Keep out of any body of water. Do not apply where runoff is likely.

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (Grus americana). Do not use this product during the period 30 days prior to arrival and 30 days after the whooping crane leaves known crane habitat.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep tightly sealed in a cool, dry place. Do not allow contact with water or acids.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

BEST AVAILABLE COPY

BONIDE ZINC PHOSPHIDE BAIT, EPA Reg. No. 4-303

ORCHARDS AND GROVES: For control of meadow mice and pine mice in orchards and groves. Apply bait after harvest while plants are in a non-bearing phase.

Hand Baiting: Near the base of each infested tree, place one teaspoon of bait at 2-4 locations, either on surface trails or at the mouth of holes leading to underground burrow systems.

* Do not allow baits to be exposed on ground. Do not disturb the runway system. Apply at the rate of 2-3 lbs./acre.

Trailbuilder: Drop one teaspoon of bait at 4-5 foot intervals in the artificial trail made by the machine just inside the drip line on both sides of the trees. Apply at the rate of 2-3 lbs./acre.

Broadcast Baiting: Broadcast evenly under infested trees by cyclone seeder, or wear rubber gloves and broadcast by hand. Concentrate in areas with the heaviest vegetation cover. Do not broadcast on bare ground or growing crops. Apply at the rate of 6-10 lbs./acre.

Aerial Application: Apply at the rate of 6-10 lbs./acre. Do not apply on bare ground or growing crops.

ORNAMENTAL AND NURSERY STOCK (Such as forest nurseries, Christmas tree nurseries, forage crops, nurseries and landscape plantings): For control of meadow mice and pine mice in ornamental and non-bearing nursery stock. Do not apply on roads, over water, or where plants are grown for food or feed.

Hand Baiting: Near the base of infested stock, place one teaspoon of bait at 2-4 locations, either on surface trails or at the mouth of holes leading to underground burrow systems. Do not allow baits to be exposed on ground. Do not disturb the runway system. Apply at the rate of 2-3 lbs./acre.

Broadcast Baiting: Broadcast evenly under infested nursery stock or non-bearing fruit trees by cyclone seeder, or wear rubber gloves and broadcast by hand. Concentrate in areas with the heaviest vegetation cover. Do not broadcast on bare ground. Apply at the rate of 6-10 lbs./acre.

HAND BAITING IN NON-CROP AREAS (Such as lawns, ornamentals, golf courses, parks and nurseries): USE RESTRICTIONS: For control of meadow and pine mice (voles, *Microtus*, spp.), Thirteen-lined ground squirrels (*Spermophilus tridecemlineatus*), Columbian ground squirrels (*Spermophilus columbianus*) and Richardson's ground squirrels (*Spermophilus richardsonii*). This product must not be applied on roads, over water or where plants are grown for food or feed.

* Control artificial trails (eg in nuts, woods, or jungles) or by pulling overhanging grass back to place.

NOT AVAILABLE COPY

BONIDE ZINC PHOSPHIDE BAIT, EPA Reg. No. 4-303

Prebaiting with 6 pounds of untreated oat groats per acre, 2-3 days prior to applying toxic bait has been shown to provide more consistent acceptance by target species when baiting with acute toxicants.

HAND BAITING: Place 4 grams (one teaspoon) of bait around each active burrow or runway, allow bait to fall through grass to ground surface but do not apply to bare ground. Do not place in heaps or piles.

BUILDINGS (homes, industrial agricultural and commercial buildings, and similar structures): For control of Norway rats, roof rats and house mice. Bait must be placed in locations not accessible to children, pets, domestic animals and wildlife, or in tamperproof bait boxes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

General Information: Remove as much food as possible and determine dry, acid-free areas where rodents are likely to find and consume the bait (usually along walls, gnawed openings, or beside burrows, in corners and concealed places, between floors and walls, and wherever evidence of rodents has been observed). Provide a continuous supply of fresh bait until all signs of feeding have stopped. Do not treat the same area with bait at less than 30 day intervals. Collect and dispose of all dead animals and excess bait properly.

Norway Rats and Roof Rats: Apply 1 or 2 teaspoons of bait per placement. Outdoors - place bait in active rat burrows or tamperproof boxes.

House Mice: Apply 1 teaspoon of bait per placement at 8-12 ft. intervals. Outdoor - protect bait from rain or snow.

Guarantee limited to label claims.

Mfg. by BONIDE CHEMICAL CO., INC., YORKVILLE, NY 13495

BEST AVAILABLE COPY