

BONIDE

Rose, Flower & Ornamental
Insect Spray
READY-TO-USE

AN EFFECTIVE INSECTICIDE FOR USE ON
ORNAMENTAL PESTS. KILLS GYPSY MOTH
CATERPILLARS AND JAPANESE BEETLES IN
ADDITION TO OTHER LISTED INSECTS.

ACTIVE INGREDIENTS	
*Pyrethrins.....	0.02%
**Piperonyl Butoxide, Technical.....	0.20%
INERT INGREDIENTS.....	99.78%
	100.00%

*From PYRENONE - Registered trademark of Agrevo
**Equivalent to min. 0.16% (butylcarbityl) (6-propyl-piperonyl) ether and 0.04% related compounds.

CAUTION - KEEP OUT OF REACH OF CHILDREN
See back panel for additional Precautionary Statements.

DO NOT TRANSPORT OR STORE UNDER 32°F
If freezing occurs, thaw and agitate vigorously.

NET CONTENTS: _____

ACCEPTED
APR -7 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 4-309

BEST AVAILABLE COPY

272


PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS-CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.



ACCEPTED

APR - 7 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 4-309

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Do not store under 32 F. Do not reuse empty container. Wrap container and put in trash collection.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

To Operate: Shake Before Using: Turn nozzle of sprayer to "spray" or "stream" position. Prime sprayer by holding at cap level of bottle and pumping trigger. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage. Set nozzle to closed position.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season. Repeat treatment weekly or as needed to control infestations.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladioli and Marigolds. **To control:** Aphids, flea beetles, leafhoppers and whiteflies. **To kill:** Gypsy moth caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

FOR USE ON VEGETABLES TO CONTROL INSECTS

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Kohlrabi, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes. **To Control:** Asparagus beetles, Aphids, Leafhoppers, Whiteflies, Mexican bean beetles, 12-Spotted cucumber beetles, Cabbage loopers, Cross-striped cabbageworms, diamondback moth larvae, Flea beetles, Harlequin bugs, Imported cabbageworms, Stink bugs, Green peach aphids, Leafhoppers, Fireworms, Blister beetles, Colorado potato beetles and Webworms.

FOR USE AS A FLYING INSECT SPRAY

Indoors as a Space Spray: To control Houseflies, Fruit flies, Mosquitoes, Gnats and Small flying moths: Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct mist toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

Outdoors, Adult Mosquito Control: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest, particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at the rate of 4 ounces per 1,000 square feet (5 to 6 quarts per acre).

To control mosquito larvae, apply at the rate of 23 ounces per 1,000 square feet (7.75 gallons per acre). Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

Buyer assumes all risks of use, storage or handling of this products not in strict accordance with directions given herewith.

Manufactured by BONIDE Products, Inc. - Yorkville, NY 13495

BEST AVAILABLE COPY