



DAIRY & HORSE FLY SPRAY

[Additional Brand Name: Pony, Horse & Pet Spray]

Optional Marketing Statements:

H₂O Water Based

Kills and Repels

A PYRENONE[®] Micro-Emulsion – Clear – Stable

Leaves no oily or objectionable residues

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants, Dairies, on Livestock and Outdoors for Mosquito Control

For Control of Fleas and Brown Dog Ticks on Premises and on Pets

Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS:

Pyrethrins (CAS #8003-34-7)..... 0.1%

*Piperonyl Butoxide, Technical (CAS #51-03-6) 1.0%

OTHER INGREDIENTS:..... 98.9%

TOTAL..... 100.0%

*Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE[®] – Registered Trademark of Bayer AG.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back/side/other panel(s) for additional precautionary statements.

ACCEPTED
 DEC 1 2003
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. **4-331**

DO NOT TRANSPORT OR STORE UNDER 32°F.

NET CONTENTS: _____

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency medical treatment information.	
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before space spraying or surface applications.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
<p>Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.</p> <p>Disposal: <i>If empty</i> – Do not reuse this container. Place in trash or offer for recycling if available. <i>If partly filled</i> – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>

SHAKE WELL BEFORE USE

LIVESTOCK: To protect cattle, horses and hogs from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair

thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

TO CONTROL POULTRY LICE: spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES: in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS & CATS. Do not repeat treatment for 7 days. Do not use on puppies or kittens under 12 weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals.

ON PETS: For fleas and ticks, start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

In case of Emergency, call 1-800-548-2423 (Animal Poison Control Center).

PET AREAS: To control Fleas and Brown Dog Ticks on premises: Turn sprayer nozzle to spray position. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

INDOOR APPLICATION:

Cover all food utensils, fish tanks, and pet food before use. Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

SURFACE SPRAYING to Control Crawling Insects: Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clover mites, Cheese Mites, Earwigs, Sowbugs, Boxelder Bugs, Centipedes, Grainary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Mealworms, Grain Mites and Cadelles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

To control **carpet beetles**: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

To control **bedbugs**: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control **fleas and brown dog ticks**: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

To control **clothes moths**: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

Space Spraying: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths**: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3 – 6 ounces per 1000 cubic feet of space.

Fogging: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet (1 qt. to 5,000 to 10,000 cubic feet) of space. Contact insects directly whenever possible.

Flying Insects: Flies, Houseflies, Mosquitoes, Gnats, Mediterranean Flour Moths, Small Flying Moths. Turn sprayer nozzle to spray position. Keep about three feet away from walls, fabrics and furniture.

Wasps & Hornets: Turn sprayer nozzle to stream position. Spray directly on insects whenever possible. Apply a wet spray after dark when all insects have returned to their nests.

Garden Insects on House Plants and Ornamental Plants Outdoors: Turn sprayer nozzle to spray position. Direct spray so that upper and lower leaf surfaces are contacted. To control Aphids, Bagworms, Thrips, Flea Beetles, Japanese Beetles, Lace Bugs, Leafhoppers, Leaf tiers, Mimosa Webworm Larvae, Plant Bugs and Whiteflies use fine mist spray about 18 to 24 inches from infested plants. Use caution on new growth and tender foliage. Repeat treatment as necessary. Avoid spraying plants when temperature exceeds 85°F. Spray when air is still. Avoid heavy wetting.

Adult Mosquito Control: Apply as a fine mist to thoroughly treat the entire area. For best result, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest (low marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 1000 square feet, 1 oz. per 10,000 sq. ft.) Using truck-mounted conventional misting equipment such as Rotomist models 51C.91 or 100HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 0.9 gallon per minute.

As an Insect Repellent: (Camping, fishing, hunting) – Spray outer clothing and surroundings. Repels Biting Flies, Mosquitoes, Midges and Gnats. If sprayed every hour or so, will aid in keeping surroundings free from Mosquitoes. If in rare cases skin irritation results, avoid continued contact. Do not pour or squirt – use a good sprayer which will throw a heavy fine mist.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents, call the National Pesticide Telecommunications Network at 1-800-858-7378.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Manufactured by:
Bonide Products, Inc.
Oriskany, NY 13424