

4-333

02/16/2001

31A 4-11/3

Instructions on reverse before complete

Form Approved

Form Approved

No. 2070-0060, Approval expires 2-28-95



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 4-333
2. EPA Product Manager George Tompkins
3. Proposed Classification None Restricted
4. Company/Product (Name) Bonide Slug, Snail & Sowbug
5. Name and Address of Applicant (Include ZIP Code) Bonide Products, Inc. 6301 Sutliff Rd. Oriskany, NY 13424
6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name

X Check if this is a new address

Section - II

Amendment - Explain below. Resubmission in response to Agency letter dated. Notification - Explain below. Final printed labels in response to Agency letter dated. "Me Too" Application. Other - Explain below. FEB 16 2001

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

No Escape Slug, Snail & Sowbug Bait

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations of 40 CFR 152.46, and no other changes have been made to the labeling or the CSF of this product. Bonide Products, Inc. understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA and further understands that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and Bonide Products, Inc. may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged in: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging, 2. Type of Container, 3. Location of Net Contents Information, 4. Size(s) Retail Container, 5. Location of Label Directions, 6. Manner in Which Label is Affixed to Product

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name: Cheryl L. Schultz, Title: State Registration Manager, Telephone No. (315) 436-8231

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received

Received

(Stamped)

Grid for date stamp

Signature: Cheryl L. Schultz, Typed Name: Cheryl L. Schultz

3. Title: State Registration Manager, 5. Date: 2/2/01

Slugs, Snails, Ants, Earwigs, Cutworms, Armyworm
Sowbugs, Millipedes, Crickets, Mole Crickets, Grasshoppers.

PROTECTS: Flowers, Ornamentals, Shrubs, Lawns and cer-
tain listed vegetables. Remains effective after rain or sprinkling.

HOW TO USE: Be sure area to be baited is damp. In dry
weather, water area well before baiting. Apply bait in evening
or at night as these pests feed mainly at night. Until under
control, bait should be renewed every two weeks or more
often for heavy infestations. Slugs and snails favor dark,
damp places. Bait especially well shaded areas around flow-
ers, shrubs, rockeries and other ornamentals where these
pests stay during daylight hours. Slugs, snails and the listed
insects are attracted to the bait. Place bait upwind to the
infestation for best results. Upwind is the direction from which
the wind is blowing from bait toward pests.

Slugs, snails, and the listed insects travel or migrate exten-
sively. When invasions from adjacent areas are a probability,
a lightly applied band of bait the entire length of the infested
area will provide a barrier and aid in control.

FLOWERS, ORNAMENTALS, SHRUBS, GROUND COVERS -
To kill slugs, snails, armyworms, earwigs, sowbugs, millipedes,
crickets, chinch bugs and grasshoppers broadcast the bait light-
ly over the infested area using 1 pound per 250 square feet.

DICHONDRA LAWNS, TURF LAWNS - To kill slugs, snails, and
mole crickets, water infested areas thoroughly. Broadcast direct-
ly on lawn or turf. Do not water treated areas for 48 hours. Use
1 pound to cover 1,500 sq. ft. of infested area.

VEGETABLES - To kill slugs, snails, armyworms, cutworms,
earwigs, sowbugs, crickets, mole crickets, grasshoppers, broad-
cast lightly on a soil surface around plants using rates as noted.
Do not allow bait to contact foliage or edible part of crops.
Subsequent planting of food and feed crops in treated soil must
be made with crops on this or other carbaryl labels.

Buyer's Guarantee Limited to Label Claims.
BONIDE PRODUCTS, INC.
ORISKANY, NY 13424



NOTIFICATION

FEB 16 2001

TRUSTED SINCE 1926

BONIDE

Can be used
around vegetables
gardens

NO ESCAPE

Slug, Snail & Sowbug Bait

R · e · a · d · y t · o U · s · e

This exceptional apple pumace bait
will attract and kill slugs, snails, ants,
earwigs, cutworms, sowbugs,
millipedes, crickets, grasshoppers,
armyworms and chinchbugs around
flowers, shrubs, lawns, and certain
listed vegetables. Read entire label
before use.

BONIDE NO ESC 3LUG, SNAIL & SOWBUG BAIT
EPA Reg. No. 4-33

PRECAUTIONARY STATEMENTS — HAZARDS TO HUMANS AND DOMESTIC ANIMALS — CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing.

USER SAFETY REQUIREMENTS:

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use. Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet. Do not allow children or pets to contact treated area until dust has settled.

IMPORTANT: Keep away from children, dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

FIRST AID:

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat.

If in eyes: Flush with plenty of water. Get immediate attention. **If on skin:** Remove contaminated clothing and wash affected areas with soap and water.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

Wearing household latex or rubber gloves, apply bait with a shaker can or broadcast bait as appropriate. Apply a thin even layer of bait to application area.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with treated surfaces. Allow dust to settle in treated areas before reentering. Do not repeat application more than once a week.

SLUG, SNAIL & SOWBUG BAIT Ready to Use	
Observe the number of days before harvest Preharvest Interval (PHI) for last application as noted in chart:	
Crop	Days between final application and harvest
Asparagus	Application to spears: Minimum single application rate of .9 lb./1,000 sq. ft. application with a maximum seasonal application rate of 2.7 lb./1,000 sq. ft./crop. Application to ferns (after harvest of spears): minimum single application rate of .9 lb./1,000 sq. ft./application with a maximum seasonal rate of 1.8 lb./1,000 sq. ft./crop. Retreatment interval 7 days when applied to ferns after harvest. Retreatment interval - 3 days when applications are to spears. PHI = 1 day. The total maximum seasonal application rate for spears and ferns is 4.5 lb. per 1,000 sq. ft./crop. The total number of applications is 3 prior to harvest with 2 additional applications to ferns only.
Beans, fresh and dried (snap, navy, and kidney); peas, fresh and dried; cow-pea, lentil	Minimum single application rate of .65 lb./1,000 sq. ft. application with a maximum seasonal application rate of 2.7 lb./1,000 sq. ft./crop Retreatment interval - 7 days PHI = 3 days (succulents) PHI = 21 days (dried) PHI = 14 days (storage) PHI = 21 days (hay)

↑
Lift & Peel Here