

4-341

2-8-1988

1/3

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 4-341	DATE OF ISSUANCE FEB - 8 1988
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	NAME OF PESTICIDE PRODUCT Bonide Disease Beater with Bayleton .5%	
	NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="text-align: right; margin-right: 50px;"> ASG <u>195913</u> 160 18 </div>	
Bonide Chemical Company, Inc. 2 Wurz Avenue Yorkville, NY 13495		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name for its use if it has been covered by others.		
<p style="text-align: center;">This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data. 2. Add the phrase "EPA Registration No. 4-341" to your label before you release the product for shipment. 3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(a). Your release for shipment of the product constitutes acceptance of these conditions.</p> <p>A stamped copy of the label is enclosed for your records.</p>		
Lois A. Rossi Product Manager (21) Fungicide-Herbicide Branch Registration Division (TS-767C)		
Enclosures		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE

BONIDE

2/3

X - GREEN

TREATS UP TO 2,500 SQ. FT.
NET WEIGHT 7 1/2 LBS.
EASY TO USE GRANULES
FOR CONVENIENT SPREADER APPLICATION

ACCEPTED
FEB - 8 1988
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. 4-341

DISEASE BEATER

Lawn Fungicide

with

• **BAYLETON Fungicide**

pink

X - yellow

controls

X - GREEN

**DOLLAR SPOT, COPPER SPOT, RED THREAD,
STRIPE SMUT, FUSARIUM BLIGHT, ANTHRACNOSE, RUSTS,
PINK AND GRAY SNOW MOLD, SOUTHERN BLIGHT**

EPA Est. # NY-1

EPA Reg. No. 4-341

ACTIVE INGREDIENT:

1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone

INERT INGREDIENTS:

BAYLETON is a Reg. TM & Serv. Ad. Lawn Fungicide

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

Note back panel for Statements of Practice, Treatment Directions, and Precautions

05
995
1000

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

STATEMENTS OF PRACTICAL TREATMENT

If **swallowed**, administer water freely and induce vomiting by giving one dose (½ oz. or 15 ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

If **on skin**, wash skin immediately with soap and warm water. If irritation occurs, get medical attention.

If **eyes are contaminated**, wash with plenty of water for at least 15 minutes. Get medical attention.

If **inhaled**, remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

In case of poisoning, call a physician or Poison Control Center.

ENVIRONMENTAL HAZARDS

Do not use on crops grown for food or forage. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not make applications when weather conditions favor drift from target area.

when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions before using.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

Container Disposal: Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed above. In spill or leak incidents, keep unauthorized people away. You may contact Chemtrec at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

AVISO PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(**TO THE USER:** If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

APPLICATION

BAYLETON is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control.

NOTE: Apply the specified amount of Bonide Disease Beater with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

RECOMMENDED APPLICATIONS • TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of Bonide Disease Beater Per 1,000 Sq. Ft.		REMARKS
	Preventative	Curative	
Dollar Spot	3	—	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	1.5	—	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
	—	6	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.
Copper Spot Powdery Mildew Red Thread (<i>Laetisaria luciformis</i>) Rusts Brown Patch (<i>Rhizoctonia solani</i>) (Suppression)	3	6	PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use a longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions. CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.
Anthracnose Poa Annual Decline	6	—	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	3	—	PREVENTIVE RATE: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	—	6	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.
Stripe Smut	Preventive Rates Only 6		Make a single application in spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phaeocephala graminicola</i>)	6		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Gray Snow Mold (<i>Typhula incarnata</i>)	12		Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold Fusarium Patch (<i>Fusarium nivale</i>) Southern Blight (<i>Sclerotium rolfsii</i>)	6 to 12		Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rates in areas with a history of severe disease damage.

Buyer's Guarantee Limited to Label Claims

MFG.
~~Sold by~~

3
B