

4-342

8/26/2003

1/4

ACCEPTED
AUG 26 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-342



LAWN FUNGICIDE WITH BAYLETON® FUNGICIDE

Optional Marketing Statements:

Treats up to 10,000 sq. ft. [15 lb. package]
 Easy-to-use granules for convenient spreader application
 Controls...Dollar Spot, Copper Spot, Red Thread, Stripe Smut, Fusarium Blight, Anthracnose,
 Rusts, Pink and Gray Snow Mold, Southern Blight.

ACTIVE INGREDIENT:

Triadimefon (CAS # 43121-43-3): 1%

OTHER INGREDIENTS:..... 99%

TOTAL..... 100%

Bayleton is a Reg. TM of Bayer AG, Leverkusen.

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

Note back/side/other panel(s) for First Aid and additional precautionary statements.

NET WEIGHT: _____

BONIDE LAWN FUNGICIDE with BAYLETON® Fungicide,

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency medical treatment information.	
If Swallowed:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.
Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

ENVIRONMENTAL HAZARDS

Do not use on crops grown for food or forage. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.
Do not make applications when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions before using.

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Disposal: *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

APPLICATION

BAYLETON® is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control.

NOTE: Apply the specified amount of Bonide Lawn Fungicide with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of ¼ inch of water has been applied to treated area. Do not use clippings for animal feed.

RECOMMENDED APPLICATIONS · TURF GRASS DISEASE CONTROL

DISEASE	Lbs. of Bonide Lawn Fungicide Per 1,000 Sq. Ft.		REMARKS
	Preventive	Curative	
Dollar Spot	1.5	-	PREVENTIVE RATE: Apply recommended rate on 30 day intervals. Protective activity of BAYLETON may extend for as long as 60 days dependent upon environmental conditions.
	0.75	-	PREVENTIVE RATE: Apply recommended rate on 14 day intervals. Protective activity of BAYLETON may extend for as long as 30 days dependent upon environmental conditions.
		3	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.
Copper Spot Powdery Mildew Red Thread <i>(Laetisaria fuciformis)</i> Rusts Brown Patch <i>(Rhizoctoni solani)</i> (Suppression)	1.5		PREVENTIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light to moderate disease development, use longer interval. Protective activity of BAYLETON can be greater than 30 days depending on environmental conditions.
		3	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.

Anthracnose Poa Annua Decline	3	-	PREVENTIVE RATE: Apply at 30 day intervals and repeat as necessary for seasonal control. Dependent upon environment conditions, residual control may be extended to 45 days.
	1.5	-	PREVENTIVE RATE: Apply at 21 day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 30 days.
	-	3	CURATIVE RATE: To control existing infections, apply the curative rate. Subsequent applications should be applied at a preventive schedule and rate.
Stripe Smut	Preventive Rates Only 3		Make single application in spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phialophora graminicola</i>)		3	Apply first application in mid - June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.
Gray Snow Mold (<i>Typhula incarnata</i>)		6	Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mold Fusarium Patch (<i>Fusarium nivale</i>) Southern Blight (<i>Sclerotium rolfsii</i>)		3 to 6	Apply before conditions favorable to infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.

Buyer's Guarantee Limited to Label Claims

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents, call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by:

Bonide Products, Inc.

Oriskany, NY 13424