

100110


Blowflies, Flies, Mosquitoes, Hcg Lice, Ticks, including Deer Ticks	Dilute 8oz in 33 gal water Dilute 1pt in 66 gal water Dilute 1qt in 132 gal water	Spray, dip or sponge to apply 1 pt per pig, especially around ears.
Poultry Mites, Northern Fowl Mites, and Lice	Dilute 8oz in 16.5 gal water Dilute 1pt in 33 gal water Dilute 1qt in 66 gal water	Spray at the rate of 1/2 per bird, or 1gal per 100 birds, with a fine mist. Spray roosts, walls and nests or cages.

**PESTS ON DOGS (DO NOT TREAT PUPPIES LESS THAN 4 WEEKS OLD)**

Fleas, Ticks, including Deer tick, Lice and Mange	Dilute 8oz in 16.5 gal water Dilute 1pt in 33 gal water Dilute 1qt in 66 gal water	Thoroughly wet the an- imal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Gives long lasting pro- tection against re- infestation and ap- plications at intervals of 2 to 3 weeks. Repeat applications as necessary.
---	--	---

**WARRANTY STATEMENT**

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.



Packed By  
BONIDE PRODUCTS, INC. - 2 WURZ AVE - YORKVILLE, NY 13495

PM13 4-358 10/8

US ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (75-767)  
WASHINGTON, DC 20460

EPA REGISTRATION NO.

4-358

DATE OF ISSUANCE

MAY 10 1993

TERM OF ISSUANCE

Until Reregistration

NAME OF PESTICIDE PRODUCT

Bonide RTU Permethrin Spray

NOTICE OF PESTICIDE:  REGISTRATION  
 REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Bonide Products Inc.  
2 Wurz Ave.  
Yorkville, NY 13495

~~160543779L~~  
18/1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) since you have agree to:

1. Submit and/or cite all data or other material for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phrase, "EPA Registration No. 4-358" before you release the product for shipment.
  - b. The product name is misleading since the label provides for multiple uses of this product for indoor and outdoor pest control. Thus, revise the product name so it is not specific to horses or else limit use of the product to horses.
  - c. You indicate this product is ready to use. How does one apply the product? Is there a spray head attachment that fits on the container head? Products intended for homeowner use should be marketed with some sort of pump spray attachment or reference should be made on the label informing the user of certain pump spray trigger heads that can be used with the product.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

2018

In addition, a label statement should be added which directs the user to attach the spray head if it is not already affixed to the container, i.e.,:

Affix trigger, pump [or other type] sprayer to container for application. Do not transfer to another container.

3. Submit 5 copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Enclosures

FRONT PANEL

BONIDE

**RTU HORSE & STABLE PERMETHRIN SPRAY**

EPA Est. No. 4-NY-1

EPA Reg. No. 4- \_\_\_\_\_

FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS

FOR CONTROL OF FLEAS, BROWN DOG AND DEER TICKS, AND LICE  
ON PREMISES AND DOGS

CONTROLS FLEAS AND TICKS ON DOGS FOR 35 DAYS AND  
AIDS IN CONTROL OF TICKS ON DOGS FOR AN ADDITIONAL 2 WEEKS

ACTIVE INGREDIENT:

\*Permethrin ..... 0.50%

INERT INGREDIENTS: ..... 99.50%

TOTAL 100.00%

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)  
2,2-dimethylcyclopropanecarboxylate, Cis/trans ratio: min. 35%  
(+/-) cis and max. 65% (+/-) trans.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See back panel for additional precautionary statements

NET CONTENTS 1 QT.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

MAY 10 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg.

4-358

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS - CAUTION**

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING

**INDOOR APPLICATION**

For spot application only. Do not use as a space spray indoors. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

**SURFACE SPRAYING:**

To control Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetle, Meal Worms, Grain Mites and Caddisflies: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards, moldings, and floors. Repeat treatment as needed.

To control Fleas (adults and larvae) and Ticks, including Ticks which may carry Lyme disease (adult and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

#### ON LIVESTOCK

To protect cattle (beef and dairy), goats, sheep, hogs, and horses from Horn Flies, Face Flies, House Flies, Mosquitoes, and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control Stable Flies, Horse Flies, and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

6048

To control *Bedbugs* and *Mites* in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

#### ON DOGS

To control *Fleas*, *Ticks*, and *Lice*: Start spraying at the tail moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. DO NOT USE CATS OR KITTENS. Do not use on old, sick, or debilitated animals.

#### OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):

Use Bonide RTU Horse & Stable Permethrin Spray as a residual treatment to control *Ticks* including the *Deer Tick* (*Ixodes dammini*), *Lone Star Tick* (*Amblyomma americanum*), *Brown Dog Tick* (*Rhipicephalus sanguineus*), *American Dog Tick* (*Dermacentor variabilis*) and *Western Black-legged Tick* (*Ixodes pacificus*).

For each areas listed below, apply until surface is wet (approximately 1 gal/750-1000 ft<sup>2</sup>).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around and the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 to 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**FOR CONTROL OF GARDEN INSECTS:**

For use on roses, dahlias, aster and other ornamentals to kill Japanese Beetles hit by spray as well as Whiteflies, Aphids, Exposed Thrip, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews, and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites, and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

Buyer assumes all risks of use, storage, or handling of this product not in strict accordance with directions given herewith.