



4-358 7-30-2003 1-9

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 4-358	2. EPA Product Manager LaRocca	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Bonide Permakil RTU Permethrin Spray	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Bonide Products, Inc. 6301 Sutliff Road Oriskany, NY 13424 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**NOTIFICATION**  
JUL 30 2003

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION: To bring label into compliance with PR Notices 96-6 & 2001-6.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula of this product. Bonide Products, Inc. understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA, and further understands that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and Bonide Products, Inc. may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Rebecca M. Horton	Title Consultant/Agent	Telephone No. (Include Area Code) (540) 375-8826
---------------------------	---------------------------	---

<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			8. Date Application Received (Stamped)
2. Signature <i>Rebecca M. Horton</i>	3. Title Consultant/Agent		
4. Typed Name Rebecca M. Horton	5. Date 7-18-03		

2-9

[FRONT PANEL]



# PERMAKIL RTU PERMETHRIN SPRAY

[Additional Brand Name: Bonide Permethrin Spray Ready-to-Use]

Optional Marketing Statements:

For Use on Roses, Begonias, Violets, and Other Listed Ornamental Plants

For Domestic Indoor and Outdoor Use Only

Kills Fire Ants

For Control of Fleas, Brown Dog Ticks, Deer Ticks and Lice on Premises and Dogs

Controls Fleas and Ticks on Dogs for 35 Days and Aids in Control of Ticks on Dogs for an Additional 2 Weeks

Controls Insect Pests on Vegetables and Flower Gardens

To Control Pests of Livestock

Provides 60 Day Control of Cockroaches, Fleas and Stored Product Pests

**NOTIFICATION**

**JUL 30 2003**

**ACTIVE INGREDIENT:**

Permethrin (Cas # 52645-53-1) ..... 0.50%

**OTHER INGREDIENTS:**..... 99.50%

**TOTAL** ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See back/side/other panel(s) for additional precautionary statements.

**NET CONTENTS:** \_\_\_\_\_

EPA Reg. No. 4-358  
7/18/03 – notification

EPA Est. No. 4-NY-1  
Page 1 of 7 Pages

The last digit in the lot number indicates EPA  
Establishment location.

[BACK/SIDE/OTHER PANEL(S)]

**READ ENTIRE LABEL BEFORE EACH USE.**

### **PRECAUTIONARY STATEMENTS**

#### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL BEFORE USING**

Affix trigger sprayer to container for application. Do not transfer to another container.

#### **INDOOR APPLICATION**

For spot application only. Do not use as a spray indoors. Do not apply to classrooms while in use. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

#### **SURFACE SPRAYING:**

To control Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetle, Meal Worms, Grain Mites and Cadelles:

Direct the spray into hiding places, cracks, crevices, behind shelves and drawers. For silverfish, spray bookcase. For ants, spray trails, nests and points of entry.

To control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor covering and baseboard, molding, and floor. Repeat treatment as needed.

EPA Reg. No. 4-358  
7/18/03 – notification

EPA Est. No. 4-NY-1  
Page 2 of 7 Pages

The last digit in the lot number indicates EPA  
Establishment location.

To control Fleas (adult and larvae) and Ticks, including Ticks that may carry Lyme disease (adult and larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboard, molding, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

To control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

**ON LIVESTOCK**

To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes, and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

To control Stable Flies, Horse Flies, and Deer Flies: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood-sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

To control Poultry Lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To control Bedbugs and Mites in poultry houses: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**USE ONLY ON DOGS. DO NOT USE ON CATS.** Do not repeat treatment for seven (7) days. Do not use on puppies under (12) weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals.

To control Fleas, Ticks, and Lice: Start spraying at the tail moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets

3-9

thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat as necessary.

In case of an emergency, call 1-800-548-2423 (Animal Poison Control Center).

#### OUTDOOR APPLICATION

##### TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):

Use Bonide Permakil RTU Permethrin Spray as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Dermacentor variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each area listed below apply until surface is wet (approximately 1 gal/750-1000 sq. ft.).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around and the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6 to 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**TO CONTROL FIRE ANTS:** Apply 1 gallon as a gentle rain to each fire ant mound using a sprinkler can or other applicator. Thoroughly wet the mound and surrounding area to a 4 foot diameter. For best results, apply in cool weather, 65-80 degrees F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound. Do not use equipment that produces a pressurized spray as pressurized sprays may disturb fire ants and cause migration, reducing product effectiveness.

**PESTS ON OUTSIDE SURFACES OF BUILDINGS:** Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet where pests are and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

**HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, FLYING MOTHS OUTDOORS:** For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary.

**ANTS, LONE STAR TICKS, CRICKETS, FLEAS AND EARWIGS OUTDOORS:**

For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat applications as infestations warrant and as reinfestations occur.

**WASPS, HORNETS, YELLOW JACKETS, BEES OUTDOORS:** Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

**FOR CONTROL OF GARDEN INSECTS:**

For use on roses, dahlias, aster and other ornamentals to kill JAPANESE BEETLES hit by spray as well as Whiteflies, Aphids, Exposed Thrip, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, dephiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews, and zinnia.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, Exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites, and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting a tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 degrees F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

**DIRECTIONS FOR USE ON VEGETABLES, FRUITS, AND NUT TREES:**

For backyard gardens only.

Apply 1 quart of this product to 110 square feet, (or 1 gallon to 440 sq. ft.) of surface area. Thorough coverage of all plants is important. Begin applications when insects first appear. Repeat applications as needed or as noted on the following chart:

VEGETABLE	TARGET PEST(S)	REMARKS
Asparagus (pre-harvest)	Cutworm Asparagus	Apply for uniform coverage as needed. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
(post-harvest)	Larval stages of Asparagus Beetle, Tarnished Plant Bugs, Lygus Bugs, and Adult Japanese Beetle	Treat ferns or bush growth after spear harvest when insects are present.

Celery	Vegetable Leaf Miner, Cabbage Looper, Black and Granulate Cutworm, Fall, Beet and Southern Armyworm	Apply every 3-5 days or as needed. Can be applied up to 3 days prior to harvest. Do not make more than 10 applications per season.
Cole Crops: Broccoli Brussels Sprouts Cabbage Cauliflower	Beet Armyworm, Diamond Back Moth, Imported Cabbageworm, Cabbage Aphid, Cabbage Looper	Apply every 5 days or as needed. Can be applied up to one day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts, and Broccoli and 10 applications to Cabbage.
Eggplant	Colorado Potato Beetle, Vegetable Leafminer	Apply for uniform coverage as needed. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.
Horseradish (Foliar application)	Imported Crucifer Weevil	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.
Head Lettuce only (Do not use on Leaf Lettuce)	Cabbage Looper, Alfalfa Looper, Beet Armyworm, Corn Ear Worm and Tobacco Budworm	Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.
Potatoes	Beet Armyworm, Colorado Potato Beetle, Potato Leafhopper, Potato Flea Beetle, Aster Leafhopper, Potato Aphid, Potato Tuberworm, Cabbage Looper, Cutworms, European Corn Borer, Potato Psyllid, Tarnished Plant Bug	Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest.
Peppers	Vegetable Leafminer, Cabbage Looper, Flea Beetle, Corn Earworm, Pepper Weevil, European Corn Borer	Apply as needed, but not more than 8 sprays per season. Can be applied up to 3 days prior to harvest.
Spinach	Beet Armyworm, Corn Earworm, Cabbage Looper, Cutworms, European Corn Borer, Green Cloverworm and Leafhoppers	Apply for uniform coverage when needed but not more than 7 applications per season. Can be applied up to 7 days prior to harvest.
Sweet Corn	Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm	Apply every 5 days and when needed but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
TREES (FRUIT OR NUT)	TARGET PEST(S)	REMARKS
Almonds	Navel Orange Worm, Peach Twig Borer	Apply when insects appear. Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Plum Curculio, Redbanded Leafroller, Rosyapple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique-banded Leafroller, Green Fruitworm	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
Peaches	Lesser Peachtree Borer, Peach Twig Borer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 2 applications per season.

Pears (dormant through delayed dormant)       (summer)	Pear Psylla       Pear Psylla, Green Fruitworm and Codling Moth	Apply during the dormant through delayed dormant growth period only. Add if desired 8 oz. of Bonide All Season's oil to each gallon of this product. Do not make more than 2 applications per season.  Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.
---	---	--

**STORAGE AND DISPOSAL**

**Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

**Disposal:** *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Manufactured by  
 Bonide Products, Inc.  
 Oriskany, NY 13424

July 18, 2003

CERTIFIED MAIL #7002 3150 0000 3062 0645

Mr. George LaRocca  
Document Processing Desk - (NOTIF)  
Registration Division - (7504C)  
Office of Pesticide Programs  
U.S. Environmental Protection Agency  
Ariel Rios Bldg.  
1200 Pennsylvania Ave., NW  
Washington, DC 20460

Re: Notification  
EPA Reg. No. 4-358, Bonide Permakil RTU Permethrin Spray

*Registrations  
by Design™  
Inc.*

Dear Mr. LaRocca:

On behalf of Bonide Products, Inc. of Oriskany, New York, enclosed a notification for the above-referenced pesticide product label.

The purpose of this notification is to bring the label into compliance with PR Notices 96-6 & 2001-6.

The following documents are enclosed:

1. EPA Form #8570-1, Application for Pesticide Amendment - Notification
2. A copy of the amended label with revisions highlighted in red.

I trust this will provide the information needed for you to proceed with review of this notification.

Best regards,



Rebecca M. Horton  
Consultant/Agent for  
BONIDE PRODUCTS, INC.

Enclosures

cc: MVickers/BONIDE

Member CHEMICAL PRODUCERS & DISTRIBUTORS ASSN.

Federal & State Pesticide/Fertilizer Registration Services plus typesetting & design of final product labeling

118½ E. Main Street, Suite 1, Salem, VA 24153 • (540)375-8826 • fax (540)375-8827

e-mail: registrationsbydesign@worldnet.att.net