

PM 3

4-372

8/20/98

1/4

PYRENONE GARDEN DUST Ready-To-Use

IT IS THE LEAST YOU CAN DO FOR YOUR GARDEN

- * CONTAINS PYRETHRIN
- * FOR USE IN AND AROUND THE HOUSE
- * USE IN NON-FOOD AREAS OF RESTAURANTS AND FOOD PROCESSING PLANTS
- * INDOOR/OUTDOOR
- * PROVIDES RAPID CONTROL OF INSECTS ON VEGETABLES OR ORNAMENTAL PLANTS
- * FOR USE ON ROSES & FLOWERS, FRUITS & VEGETABLES, FOUNDATIONS, PATIOS, DECKS
- * FOR CONTROL OF LICE ON POULTRY AND PET BIRDS, FLEAS AND TICKS ON PREMISES, FLEAS AND TICKS ON DOGS, CATS, GERBILS HAMSTERS AND WHITE MICE
- * ACCEPTABLE FOR USE TO THE DAY OF HARVEST

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
Piperonyl Butoxide Technical. CAS # 51-03-05	1.00%

OTHER INGREDIENTS:	98.90%
TOTAL:	100.00%

Pyrenone-Registered Trademark of AgrEvo

EPA REG. NO. 4-372

EPA EST NO. 4-NY-1

ACCEPTED

AUG 20 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-372

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautions.

NET CONTENTS: _____

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled or absorbed through skin. Avoid inhalation of Dust. Avoid contact with skin.

PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PYRENONE GARDEN DUST may be used throughout the growing season; or as a preharvest treatment-acceptable for use up to the day of harvest on edible plants when other insecticides are prohibited due to residue restrictions and use limitations.

Treat only when insects are present and repeat every 7 days as required to maintain control. Apply dust uniformly to upper and lower leaf surfaces.

For ____ oz. Duster: Shake to fluff powder. This container is a refillable puffer applicator. Open the container, aim at plants to be treated and with your fingers rapidly squeeze and release the sides of the container. Follow the instructions for each plant.

For ____ lb. Canister: Shake to fluff powder. This container is a refillable duster applicator. Shake over plants to be treated or crush the sides of the canister in and rapidly squeeze and release in order to get a bellows dusting effect. Follow the instructions for each plant.

For ____ lb. Applicator Bags: This bag is a self contained Duster Bag. Cut off the bottom at the line indicated. Invert the bag and gently shake to move dust away from the screen. Open the Zip-Loc closure. Slowly turn the bag over and dust plants by holding the handle and gently shaking up and down. Follow the instructions for each plant. When finished, invert the bag and reseal the Zip-Loc for storage.

CROP	TO CONTROL
Asparagus Beans	Asparagus Beetles, Aphids, Leafhoppers, Mexican Bean Beetles
Broccoli, Brussel Sprouts, Cabbage, Cauliflower	Cabbage Loopers, Diamondback Moth Larvae, Flea Beetles, Harlequin Bugs, Imported Cabbageworms
Celery	Cabbage Loopers, Leaf Tiers
Lettuce	Cabbage Loopers, Diamondback Moth Larvae, Imported Cabbageworms
Potatoes	Aphids, Colorado Beetles, Flea Beetles, Leafhoppers
Radishes	Aphids, Flea Beetles
Spinach	Aphids, Cabbage Loopers, Webworms
Tomatoes	Green Peach Aphid, Flea Beetle
Ornamentals	Aphids, Flea Beetles Japanese Beetles, Leafhoppers on: African Violets, Aster, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwood, Geraniums, Gladiolus, Marigolds, Roses, etc.

Fruit:

Citrus Fruits: Including (but not limited to) Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummel, Satsuma Mandarin.

Pome Fruits: Including (but not limited to) Apple, Crabapple, Loquat Pear, Orienta.

Stone Fruits: Including (but not limited to) Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum.

Small Fruits and Berries: Including (but not limited to) Blackberry Blueberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Grape Huckleberry, Loganberry, Olallie Berry, Raspberry (black & red) Strawberry, Youngberry.

For the Control of Insects: Such as Ants, Aphids, Armyworms, Blister Beetles, Cabbage Looper, Caterpillars, Crickets, Diamondback Larvae, Fireworms, Flea Beetles, Fruit Flies, Fruittree, Leafroller, Grape Leafhopper, Green Peach Aphids, Greenhouse Thrips, Gypsy Moth (adult & larvae), Harlequin Bug, Halitosis sp., Leafhopper, Leafrollers, Leaf tiers, Lice, Psyllids, Thrips, Webworms. Whiteflies.

OUTSIDE SURFACES OF BUILDING

Use on Ants, Clover mites, Crickets, Earwigs, Millipedes, Spiders, Cockroaches, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also to outside surfaces of buildings, including porches, decks, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen.

Other Outdoor Areas

Use on Ants, Crickets, Earwigs, Fleas, Lone Star Ticks for treatment of localized infestations of these insects in areas where there are weed or bushy non-crop areas, dust infested areas thoroughly. For ants, thoroughly treat hills and runways. Repeat application no more than once per week as needed.

Wasps, Bees and Hornets: Apply dust on nest and surrounding areas. To treat nests in wall voids, drill a hole in the wall, blow in dust and reseal. It is best to treat wasp nests at night when insects are less active and have returned to the nest. Do not wear loose fitting clothing. Take necessary precautions to avoid stings. For best results, carefully dust near nest and check one or two days after treatment to ensure complete kill, Retreat if necessary. If possible, remove and destroy nest to prevent hatching of new insects. For ground nests, carefully dust entrance and surrounding areas. Retreat in one or two days if wasp activity is still present. Do not use this product to treat overhead nests.

Nuisance Pests Indoors

Application within residential buildings including homes and apartment buildings and within non-food areas of industrial, institutional and commercial buildings including hospitals, stores, manufacturing plants and warehouses.

For Spot Treatment Only: Apply as a light but uniform dust to localized areas where **Cockroaches**, (including strains resistant to certain other insecticides), **Ants, Clover Mites, Crickets, Silverfish, and Spiders** are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerator and cabinets, around plumbing and other utility installations. **For Ants:** Apply to ant trails, also around doors and windows and wherever else these pests may find entrance.

Carpenter Ants: Apply dust around doors and windows and other places where ants enter premises and where they crawl or hide. Also force dust into cracks and crevices or through openings or small newly drilled holes into wall voids where ants and/or nests are present. Use a sufficient amount to lightly cover area surfaces.

4/4

For Control of Carpet Beetles: Apply dust as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

Fleas and Ticks on Premises: Dust thoroughly all infested areas, particularly crevices in woodwork, floor and floor covering, and animal sleeping quarters.

Bedbugs: Dismantle beds completely and dust all parts thoroughly, including, mattress seams and box springs. Dust all baseboards and wall cracks thoroughly.

Lice on Poultry: Dust the powder carefully into the feathers, under the wings, along the back and around the vent. Rub into feathers of the head. Nests, roosts, and runways, porches and cages should be thoroughly dusted. Destroy infested food and cuttlebone. Repeat no more than once per week as needed.

Lice on Pet Birds: Dust the powder carefully into the feathers, under the wings, along the back and around the vent. Rub into feathers of the head. Nests, roosts, and runways, porches and cages should be thoroughly dusted. Destroy infested food and cuttlebone. Repeat no more than once per week as needed.

Dogs and Cats

Read Entire Label Before Each Use. Do not repeat treatment for seven (7) days. Do not use on puppies or kittens under twelve (12) weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

In case of emergency, call 1-800-548-2423 (Animal Poison Control Center)

Fleas and Brown Dog Ticks on Cats and Dogs: Dust the powder onto the animal's hair and work it down to the skin. After treatment, brush or comb the animal's coat to remove dead fleas and lice. Repeat no more than once a week as needed.

Fleas on Gerbils, Hamsters and White Mice: Do not use if female is pregnant or nursing young. Dust the powder onto the animal's hair and work it down to the skin. After treatment, brush or comb the animal's coat to remove dead fleas and lice. Repeat no more than once per week as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open burning and dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by federal, state or local disposal authorities. Consult federal state or local disposal authorities for approved procedures.

For information on pesticide products (including Health Concerns, Medical Emergencies, or Pesticide Incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Buyer's Guarantee limited to label claims.

**Bonide Products, Inc.
Yorkville, NY 13495**