


PM - 10

Reg # 4-375



194

BONIDE

✓ PORCH, PATIO, GARDEN AND ORNAMENTAL SPRAY

FOR HOUSEHOLD USE ONLY

✓ KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES

A FAST ACTING EFFECTIVE WAY TO KILL WASPS AND HORNETS, MOSQUITOES, GNATS, MOTHS AND HOUSEFLIES; SPIDERS, ANTS AND ROACHES, APHIDS, GYPSY MOTH CATERPILLARS, JAPANESE BEETLES AND MORE THAN 30 OTHER YARD AND GARDEN INSECTS.

USE OUTDOORS ON FLOWERS, ROSES, TREES, AND SHRUBS, ORNAMENTALS, AND VEGETABLES - AND AROUND PORCH, PATIO AND GARAGE.

PROVIDES EFFECTIVE, FAST-ACTING, KILLING POWER WITH NO OBJECTIONABLE RESIDUE.

CAN BE USED ON VEGETABLES TILL DAY OF HARVEST

READY-TO-USE, NO MIXING NEEDED

CONTAINS NATURAL PYRETHRUM

EPA Est. No.

EPA Reg. No.

ACTIVE INGREDIENTS:

Pyrethrins

0.04%

*Piperonyl Butoxide, Technical

0.40%

**INERT INGREDIENTS:

99.56%

100.00%

*Equivalent to min. 0.32% (butylcarbityl) (6-propylpiperonyl) ether and 0.08% related compounds

See back/side panel for additional precautionary statements

Keep Out of Reach of Children

CAUTION

Harmful if swallowed

BEST COPY AVAILABLE

NET CONTENTS

2941

PHYSICAL OR CHEMICAL HAZARDS: Do not apply this product in or on electrical equipment due to the possibility of a shock hazard.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

STATEMENT OF PRACTICAL TREATMENT

Harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. If swallowed, induce vomiting by touching finger to back of throat. Contact a physician or Poison Control Center immediately. Remove pets, from areas being treated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

DO NOT TRANSPORT OR STORE UNDER 32°F

If freezing occurs, thaw and agitate vigorously.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent where its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 21 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

TO OPERATE PUMP-N-Shoot SPRAY BOTTLE: SHAKE BEFORE USING. Remove cap. Fill sprayer bottle with product to within one inch of top. Replace the cap. Pump the bottle 4 or 5 times by alternately pulling the cap up and pushing it down. Once the spray bottle is pumped up, follow label instruction for application directions. Once pumped up, the spray bottle will shoot a straight stream up to 20 feet. Spray water through the sprayer after each use. The sprayer should be rinsed thoroughly and pumped empty for storage.

BEST COPY AVAILABLE

394

VEGETABLES

Use outdoors throughout the growing season. Apply spray thoroughly on upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied to day of harvest.

INSECT CONTROLLED: Aphids, Asparagus Beetles, blister beetles, Cabbage Loopers, Colorado Potato Beetles, Cross-Striped Cabbageworms, Diamondback caterpillars, Flea Beetles, Green Peach Aphids, Harlequin Bugs, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Leftiers, Mexican Bean Beetles, Stink Bugs, 12 Spotted Cucumber Beetles, Webworms, and Whiteflies.

CROPS: Asparagus, Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collards, Cucumbers, Eggplant, Kale, Lettuce, Melons, Mustard Greens, Peppers, Potatoes, Radishes, Spinach, Squash, Sweet Corn, Tomatoes and Turnips.

ORNAMENTAL, TREES, SHRUBS AND ROSES Apply as a fine mist spraying plants on the upper and lower leaf surfaces.

INSECTS CONTROLLED: Aphids (red, black and green peach), Bagworms, Thrips, Flea Beetles, Lace Bugs, Violets, Leaf Rollers, Leaf tiers, Mimosa Webworm Larvae, Plant Bugs and Whiteflies. Repeat treatment weekly, or as needed to control infestations.

PORCHES, PATIOS AND GARAGES Spray the surface areas and hit the insects whenever possible. Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat if necessary. Wasps and Hornets: Apply in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat if necessary. Spiders: Spray directly on spiders and web. Repeat if necessary.

To kill Cockroaches, Paimetto Bugs, Earwigs, Sowbugs, Centipedes, Millipedes, Crickets, Silverfish and Clovermites. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes, spigots, storage areas, and sheds. Spray directly on insects where possible. Repeat treatment if necessary.

OUTDOORS; To Kill Mosquitoes, Gnats (midges), Houseflies, Biting Flies, and Small Flying Moths (millers), apply spray to shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion while moving away from the treated area. To reduce annoyance from the treated area. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. For best results spray when air is still. Allow spray to dissipate before occupying the treated area. Repeat if necessary. To kill Gypsy Moth Caterpillars and adults, Adult Japanese Beetles and Eastern Tent Caterpillars on contact.

On trees and shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

BEST COPY AVAILABLE

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

BUYER'S GUARANTEE LIMITED TO LABEL CLAIMS

MANUFACTURED by

BONIDE PRODUCTS, INC.

YORKVILLE, NY 13495

474

BEST COPY AVAILABLE

