



# Pyrenone Mosquito Fogging Spray

[Additional Brand Name: BONIDE® Vegetable/Ornamental Fogging Spray]

- **Ready-to-Use Oil Base Pyrethrins Insecticide Designed for Outdoor Fogging**
- **Acceptable for Indoor Use**

## ACTIVE INGREDIENTS

Pyrethrins.....	0.25%
Piperonyl Butoxide, Technical*.....	2.00%
<b>OTHER INGREDIENTS**.....</b>	<b>97.75%</b>
<b>TOTAL.....</b>	<b>100.00%</b>

\*Equivalent to 1.6% (butylcurbitlyl)(6-propylpiperonyl) ether and 0.4% related compounds.

\*\*Contains Petroleum Distillate.

PYRENONE- Registered Trademark of Roussel Uclaf Corporation


EPA Reg. No. 4-376-AMENDMENT

EPA Est. No. 4-NY-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Back/Side/Other Panel(s) for Additional Precautionary Statements

NET CONTENTS:



### FIRST AID

Call a poison control center or doctor immediately for treatment advice.

If swallowed:	Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give <b>any</b> liquid to the person. Do not give anything by mouth to an unconscious person.
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on pesticide products (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

**Note to Physician:** Contains petroleum distillate - vomiting may cause aspiration pneumonia.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Remove pets and cover aquaria before home use. This product should only be used in food processing plants when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment and washed with an effective cleaning compound followed by a potable water rinse before using.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Use only as directed and do not contaminate lakes, streams, or ponds by cleaning of equipment or disposal of wastes. Do not apply when conditions favor drift from intended area of treatment.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

**OUTDOOR MOSQUITO CONTROL:** To control adult mosquitoes apply as a fog or fine mist using good mechanical, thermal or compressed air equipment. For best results, treat when wind velocity does not exceed 5 miles per hour. Application in the evening or early morning is recommended.

Treat area thoroughly with particular attention to low and marshy places, shorelines, stagnant pools or ponds and shrubbery or vegetation where mosquitoes may rest. Apply at a rate of ~~20 fluid ounces per acre~~ (or 0.5 fluid ounces per 1000 square feet.

~~Using truck mounted or mobile equipment, the above application rate will result if a 300 foot swath is laid down traveling at 5 m.p.h. and dispersing spray at a rate of 28 gallons per hour. May also be applied by aircraft.~~

**INDOOR USE:** To control **MOSQUITOES, HOUSEFLIES, GNATS, and SMALL FLYING MOTHS.** Fill the room with mist, using one ounce per 1000 cubic feet of space. Keep room closed for 10 minutes. Vacate during treatment and ventilate before reoccupying.

**USE ON GARDENS & ORNAMENTALS:**

This insecticide may be used outdoors in gardens or in greenhouses throughout the growing season or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

For best results, use throughout the growing season to prevent buildups of insect infestations. Using good mechanical, thermal or compressed air equipment, spray plants from a distance of 1 ½ to 2 feet using 7½ - 8 fl. oz. to cover 1000 sq. ft. To prevent injury to plants, avoid visibly wetting the foliage.

Apply in the early morning or late evening when air is still, treating all foliage with particular attention to the underside of leaves. Contact insects directly whenever possible. Repeat treatment every 7-10 days or as re-infestation occurs.

Controls (Kills) most harmful garden insects, including: **Ants, Aphids, Armyworms, Asparagus Beetles, Bagworms, Beetles, Biting Flies, Blister Beetles, Cabbage Loopers, Cankerworms, Centipedes, Chiggers, Chinch Bugs, Cicadas, Clover Mites, Colorado Potato Beetles, Corn Borers, Corn Earworms, Crane Flies, Crickets, Cross-stripped Cabbage Worms, Cutworms, Diamondback Moth Caterpillars, Digger Wasps, Earwigs, Fire Ants, Firebrats, Fireworms, Flea Beetles, Fleas, Fruit Flies, Gnats, Grasshoppers, Green Bugs, Green Fruitworms,**

**Green June Beetles, Green Peach Aphids, Gypsy Moth Caterpillars and Adults, Harlequin Bugs, Hornets, Horse Flies, House Flies, Imported Cabbage Worms, Japanese Beetles, Katydid, Lace Bugs, Leafhoppers, Leafminers, Leafrollers, Leaf-tiers, Lice, Maggots, Mealybugs, Mexican Bean Beetles, Millipedes, Mites, Moths, Pillbugs, Rootworms, Silverfish, Sowbugs, Spittlebugs, Stink Bugs, Tent Caterpillars, Thrips, Ticks, Tomato Hornworms, 12-spotted Cucumber Beetles, Webworms, Weevils, Whiteflies.**

**USE ON ORNAMENTALS, ROSES & FLOWERS** including: African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, dogwood, geraniums, gladioli, marigolds, roses, rubber plants, and wandering jew.

**USE ON MOST GARDEN VEGETABLES, HERBS, FRUITS & NUTS** including: almond, apple, apricot, arrugula, asparagus, basil, beans, beet, blackberry, blackeyed peas, blueberry, broccoli, Brussels sprouts, cabbage, camomile, cantaloupe, carrot, casaba, cashew, cauliflower, celery, cherry, chicory, chives, collards, corn, cranberries, cucumbers, currant, curry leaf, eggplants, elderberry, endive, garden peas, garlic, grape, grapefruit, honeydew, horseradish, kale, kumquat, lemon, lentils, lime, lettuce, loganberry, loquat, macadamia nut, mango melon, melons, mint, mustard greens, nectarine, onion, orange, paprika, parsley, peach, pear (bitter melon), peppers, (including bell, chili), pimento, plum, potatoes, pumpkin, radishes, raspberry, sage, snap bean, soybean, spinach, strawberry, squash, sweet bay (bayleaf), sweet corn sweet potato, swiss chard, tangelo, tangerine, tomatoes, turnips, walnut, watermelon, and yam.

#### **STORAGE AND DISPOSAL**

**Storage:** Store and transport in an upright position. Store in a cool, dry area away from children and pets.

**Disposal:** *If empty* -- Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* -- Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

**Bonide Products, Inc., Oriskany, New York 13424**