PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

CAUTION: May be harmful if swallowed. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Avoid Contamination of Feed and Foodstuffs.
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste.
This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.
Do not enter treated areas without protective clothing until sprays have dried.
Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

STORAGE: Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed.
PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

SEVIN 50 W is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. Do not mix with oil base sprays.

VEGETABLES

Mix 2 tablespoonfuls per gallon of water. Apply as a full wetting spray. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary unless shorter interval is specified below.

May be used up to day of harvest on the following: BEANS: Mexican bean beetle, leafhopper, cucumber beetle, Japanese beetle. **CORN:** corn rootworm adults, European corn borer, fall armyworm, corn earworm. Applications for corn earworms should be started when silks first appear. Repeat at 2 to 3-day intervals for corn earworms or until silks are dry. **CARROTS:** flea beetles, leafhoppers, harlequin bug. **CUCUMBER, SQUASH, PUMPKINS, MELONS:** pickleworm, melonworm, cucumber beetles, squash bug, flea beetle, leafhoppers. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. **OKRA:** corn earworm, stink bug. **POTATO:** Colorado potato beetle, flea beetle, leafhopper. Apply as a foliage application only. **TOMATO, EGGPLANT, PEPPER:** tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs and lygus bugs.

Do not apply within 3 days of harvest on the following: BEETS, TURNIPS, (do not apply within 14 days of harvest if tops are to be used as food or feed.) Leafhoppers. **BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI:** flea beetle, harlequin bug, armyworm, cabbage caterpillar, and corn earworm. **RADISHES:** flea beetle, leafhopper.

Do not apply within 14 days of harvest on the following: LETTUCE (LEAF), SPINACH, SWISS CHARD: armyworm, flea beetle, leafhopper.

LAWN INSECT CONTROL

Use one level cuptul in 15 to 20 gallons of water for each 500 square feet of established lawn area for the control of: ants, chinch bugs, earwigs, fall armyworms, fleas, leafhoppers, mosquitoes, **SOD WEBWORMS** (lawn moths) millipeds. Use **GARDEN HOSE SPRAYERS** or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

ORNAMENTAL INSECT CONTROL

Ornamental plants, flowers, shrubs and shade trees including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston Ivy.

Use 2 level tablespoonfuls per gallon of spray (2 pounds per 100 gallons) when insects or their damage appear. Repeat weekly or as needed to control: apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf beetle, elm leaf aphid, flea beetle, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

SMALL FRUIT INSECT CONTROL

Use sufficient spray to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary.

GRAPES

Use 2 tablespoonfuls per 1 gallon of water for European fruit leconium, grape leaf folder, grape leafhopper, and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls, and as needed for leafhoppers. May be applied up to day of harvest.

STRAWBERRIES

For meadow spittlebug, leafroller, strawberry weevil, use 2 tablespoonfuls per gallon of water. Make 3 to 5 post bloom summer sprays as needed. May be used up to 1 day before picking.

EPA Reg. No. 3772-13
EPA Est. 407-IA-1

C-71-84-1

Manufactured For MAY SEED & NURSERY CO. Shenandoah, Iowa 51603

EARL MAY®

a amended, for the pesticide registered under EPA Reg. No. 3772-13 Transferred to 4-387

SEVIN*

WETTABLE POWDER

ACTIVE INGREDIENTS:	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	50%
INERT INGREDIENTS	50%
Total	100%

*SEVIN is a registered trade mark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 8 OUNCES

4-387

PM 19