

BONIDE

SEVIN®

brand carbaryl Insecticide

WETTABLE POWDER

	By Wt.
ACTIVE INGREDIENT:	
Carbaryl: (1-Naphthyl N-Methylcarbamate)	50%
INERT INGREDIENTS:	50%
TOTAL	100%

©SEVIN is the registered trade mark of Rhone-Poulenc for Carbaryl Insecticide.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 4-387
EPA Est. No. 4-NY-1

NET WEIGHT

ACCEPTED
FEB 10 1997
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4-387

96 NOV -5 AIO :21

RECU ... P/DPD1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Caution: May be harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. ~~Wash thoroughly after handling.~~ *Re (add) Page 3 (a.)*

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, ~~or wetlands.~~ Do not contaminate water by cleaning of equipment or disposal of waste.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Re (add) Page 3 (b.)

~~Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.~~

~~Do not enter treated areas without protective clothing until sprays have dried.~~

STORAGE: Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

SEVEN 50W is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. Do not mix with oil base sprays.

VEGETABLES

Mix 2 tablespoonfuls per gallon of water. Apply as a full wetting spray. Apply when insects or their damage appear. Repeat at 7 ~~to 10~~ day intervals or as necessary unless shorter interval is specified below. *(no more than once a week)*

May be used up to day of harvest on the following: BEANS: Mexican bean beetle, leafhopper, cucumber beetle, Japanese Beetle. CORN: corn rootworm adults, European corn borer, fall armyworm, corn earworm. Applications for corn earworms should be started when silks first appear. Repeat at ~~7 to 10~~ day intervals for corn earworms or until silks are dry. CARROTS: flea beetles, leafhoppers, harlequin bug. CUCUMBERS, SQUASH, PUMPKINS, MELONS: pickle-worm, melonworm, cucumber beetles, squash bug, flea beetle, leafhoppers. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. OKRA: corn earworm, stink bug.

POTATO: Colorado potato beetle, flea beetle, leafhopper. Apply as a foliage application only. TOMATO, EGGPLANT, PEPPER: tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bugs, lace bugs and lygus bugs.

Do not apply within 3 days of harvest on the following: BEETS, TURNIPS. (do not apply within 14 days of harvest if tops are to be used as food or feed.) Leafhoppers, BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI: flea beetle, harlequin bug, armyworm, cabbage caterpillar, and corn earworm. RADISHES: flea beetle, leafhopper. Do not apply within 14 days of harvest on the following: LETTUCE (LEAF), SPINACH, SWISS CHARD: armyworm, flea beetle, leafhopper.

GRAPES

Use 2 tablespoonfuls per 1 gallon of water for European fruit lecanium, grape leaf folder, grape leafhopper, and grape leaf skeletonizer. Apply just before first brood leaf folder larvae emerge from rolls, and as needed for leafhoppers. May be applied up to day of harvest. *(no more than once a week)*

STRAWBERRIES

For meadow spittlebug, leaf roller, strawberry weevil, use 2 tablespoonfuls per gallon of water. Make 3 to 5 post bloom summer sprays as needed. May be used up to 1 day before picking.

ORNAMENTAL INSECT CONTROL

Ornamental plants, flowers, shrubs and shade trees including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston ivy.

(no more than once a week)

Use 2 level tablespoonfuls per gallon of spray (2 pounds per 100 gallons) when insects or their damage appear. Repeat weekly or as needed, to control: apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf beetle, elm leaf aphid, flea beetle, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose-slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

SMALL FRUIT INSECT CONTROL

Use sufficient spray to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 ~~to 10~~ day intervals or as necessary. *(no more than once a week)*

LAWN INSECT CONTROL

Use one level cupful in 15 to 20 gallons of water for each 500 square feet of established lawn area for the control of: ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, SOD WEBWORMS (lawn moths) millipedes. Use GARDEN HOSE SPRAYERS or pressure type equipment and apply full water volume to insure good penetration of turf. For best results, mow lawn and apply immediately after rain or watering. Repeat 2 to 3 weeks later if necessary.

NOTICE: Buyer assumes all risk of use, storage or handling of this product when not in strict accordance with directions given herewith.

Manufactured by
BONIDE PRODUCTS, INC.
YORKVILLE, NY 13495

(a.)

USER SAFETY REQUIREMENTS

- ▶ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- ▶ Wear a hat and eye protection when making overhead applications.
- ▶ Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- ▶ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- ▶ Do not allow children or pets to contact treated area until sprays have dried.

(b.)

Apply when air is calm to avoid drift and contact with eyes and skin.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.

For trees taller than 10 feet, consider hiring a licensed professional.

Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.

Do not repeat applications more than once a week.