WTED STATE	U.S. EN	VIRONMENTAL PROTECTION Office of Pesticide Programs	AGENCY	EPA Reg. Number: 4-393	Date of Issuance:
No. 10	· ·	Registration Division (7505C) 401 "M" St., S.W.			JUL 2 1991
Real Property in the second se	• . · ·	Washington, D.C. 20460		Term of Issuance:	Conditional
THAN PROTECTION	NOTICE OF	PESTICIDE:			· .
		Registration		Name of Pesticide Pr	oduct:
(under FIFRA, as amended)		Bonide Permakil 25			
	of Registrant (include	ZIP Code):		<u> </u>	· · · · · · · · · · · · · · · · · · ·
James J. V Bonide Pro		÷ `.	· .		
2 Wurz Ave	•				
Yorkville, N	IY 13495			·	
	vision prior to use of th	stance from that accepted in po re label in commerce. In any c			
	ormation furnished by t ide and Rodenticide Ac	the registrant, the above name	f pesticide is hereby re	egistered/reregistered u	nder the Federal
the environment, t Act. The acceptar	he Administrator, on hi nce of any name in con	as an endorsement or recomm is motion, may at any time sus- nection with the registration of its use if it has been covered l	end or cancel the regi a product under this A	istration of a pesticide	in accordance with the
This pro provided th		tionally registered	in accordance	e with FIFRA	sec. 3(c)(7)(A)
under FIFR. to submit s	A sec. 3(c)(5)	l data required for when the Agency I submit acceptable A section 4.	requires all r	egistrants of a	similar products
2. Make th	ne following la	abel changes:			
a. R	evise the EPA	Registration Num	ber to read, "	EPA Reg. No.	4-393".
	-	ntence "FOR DOMI OR DOMESTIC INI			
prod	ucts. All proc	bel in accordance ducts released for s labeling that is cor	shipment by <mark>r</mark>	egistrants aft	
Signature of Appro	oving Official:	· · · · · ·	· · · · ·	Date:	
BE,	for George	Lakaca	 I		
PA Form 8570-6	<u>+ - 0</u>	<u> </u>		<u> </u>	

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Be

George T. LaRocca Product Manager 13 Insecticide Branch Registration Division (7505C)

Enclosure

# BONIDE Permakil 25

- \* FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS.
- \* A READY-TO-USE PERMANONE® STABLE EMULSION INSECTICIDE
- \* FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES AND ON DOGS
- CONTROLS FLEAS AND TICKS ON DOGS FOR 35 DAYS
- \* AIDS IN CONTROL OF TICXS ON DOGS FOR AN ADDITIONAL 2 WEEKS
- \* FOR CONTROL OF PESTS ON LIVESTOCK
- \* WITH COUNTRY FRESH FRAGRANCE
- \* PROVIDES 60-DAY (TWO MONTHS) CONTROL OF COCK-ROACHES
- \* PROVIDES 60-DAY (TWO MONTHS) CONTROL OF FLEAS
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF STORED PRODUCT PESTS

#### **ACTIVE INGREDIENT:**

•Permethrin	0.25%
INERT INGREDIENTS:	
	100.00%

\*(3-phenoxyphenył) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

PERMANONE - Registered Trademark of AgrEvo Environmental Health, Inc.

EPA REG. NO.	4- GOG	EPA EST. NO.	4 - NY - 1
--------------	--------	--------------	------------


## KEEP OUT OF REACH OF CHILDREN

## CAUTION

See Side Panel For Additional Precautions.

## NET CONTENTS:

)



### PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

#### CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

#### **Environmental Hazards**

This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING

INDOOR APPLICATION For spot application only. Do not use as a space spray indoors.

SURFACE SPRAYING: FOR 60-DAY (TWO MONTHS) CON-TROL OF: COCKROACHES, WATERBUCS, PALMETTO BUCS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUCS, PILL-BUCS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUCSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUCS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

TOCONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from *Horn Flies, Face Flies, House Flies, Masquitoes* and *Gnats*: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE: Apply to the infested areas

of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUCS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the *Bedbugs* and *Mites* hide.

ON DOGS:

)

}

TO CONTROL FLEAS, TICKS, AND LICE: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on pupplies less than four weeks old. DO NOT USE ON CATS OR KITTENS.

#### OUTDOOR APPLICATION

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill *Japanese Beetles* hit by spray as well as *Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites* and *Leafminers*. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, cameilias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, nubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur. Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

Buyer's Guarantee Limited to Label Claims

Manufactured By BONIDE PRODUCTS, INC. Yorkville, NY 13495