



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

September 30<sup>th</sup>, 2025

Rebecca Horton  
Consultant/Agent  
Bonide Products LLC  
6301 Sutliff Rd.  
Oriskany, NY 13424

Subject: Label Amendment - Registration Review Mitigation for Mecoprop-p  
Product Name: BONIDE LAWN WEED KILLER  
EPA Registration Number: 4-400  
Case Number: 477250  
Application Date: December 08, 2020

Dear Rebecca Horton:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Mecoprop-p Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact DeMariah Cotton by phone at (202)-566-2288 or via email at [cotton.demariah@epa.gov](mailto:cotton.demariah@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read 'M. Perch', with a long horizontal flourish extending to the right.

Maryam K. Muhammad-Perch, Team Lead  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

**ACCEPTED****09/30/2025**Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No.**4-400**

# LAWN WEED KILLER

[Additional Brand Names: BONIDE® Weed Beater Lawn Weed Killer; Orchard™ Est. 1931 Lawn Weed Killer Concentrate; Lawn Weed Killer Concentrate; BiMart Owner's Choice™ Lawn Weed Killer Concentrate;



Lawn Weed Killer Concentrate]

## Optional Marketing Statements:

- One Quart Covers Up to 8,000 Square Feet
- Controls Dandelion, Chickweed, Knotweed, Henbit, Plantain, Clover, Spurge, and many other broadleaf weeds including those shown on side/back panel.
- KEEP FROM FREEZING
- For Use On: Bahiagrass, Bermudagrass, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass, Centipedegrass and St. Augustinegrass
- Apply using a tank or hose-end sprayer. (see inside booklet for complete directions for use)
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Treats northern and southern grasses.
- Kills the root.
- One quart treats up to 16,000 square feet. (Southern grasses)
- Water based.
- Won't harm lawn grass when used according to label directions.
- Kills [Controls] 200+ broadleaf weeds.
- Starts working overnight
- Won't harm lawn grass [lawns] (when used as directed)
- Controls (Number) weeds!
- Mixes with Water – Concentrate
- Treated areas for this container vary from 16,000 square feet (Northern Grasses) to 32,000 square feet (Southern Grasses)

## CONTROLS

- SENSITIVE WEEDS: Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed, Sheperdspurge, and Wild Lettuce.
- HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and many other broadleaf weeds.
- For 16 fl.oz. retail container: Lawn areas for this container vary from 4,000 sq.ft.(Northern grasses) to 8,000 sq.ft.(Southern grasses).
- For 32 fl.oz. retail container: Lawn areas for this container vary from 8,000 sq.ft.(Northern grasses) to 16,000 sq.ft.(Southern grasses).
- For One Gallon retail container: Lawn areas for this container vary from 32,000 sq.ft.(Northern grasses) to 64,000 sq.ft.(Southern grasses).
- For outdoor use
- Controls dandelion, spurge, chickweed and many other hard-to control weeds.
- Contains Trimec® Herbicide

[label to feature use rate equivalents for coverage of different size containers]

## ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt .....	7.59%
Mecoprop-p, dimethylamine salt .....	1.83%
Dicamba, dimethylamine salt .....	0.84%
OTHER INGREDIENTS: .....	89.74%
TOTAL .....	100.00%

THIS PRODUCT CONTAINS:

0.54 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%

0.13 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.51%

0.06 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.70%

Isomer Specific by AOAC Method.

2,4-D, dimethylamine salt (CAS Registry No. 2008-39-1)

Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)

Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back/side/other panel(s) for additional Precautionary Statements and First Aid.

NET CONTENTS: \_\_\_\_\_

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER:</b> Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.	

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use only lawn-type sprayers. Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.
- Schedule the herbicide applications when air temperatures are 60° to 90°F with good soil moisture. Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures. The tolerance of warm season grasses decreases at air temperatures greater than 90°F and when turfgrass is drought stressed.
- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

### USE RESTRICTIONS:

Broadcast Applications - Do not exceed 17 fl. oz. per 1000 sq. ft. per application [equivalent to 0.75 lb ae mecoprop-p/A]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.

Spot Treatments - Do not exceed 27.5 fl. oz. per 1000 sq. ft. per application [equivalent to 1.2 lbs. ae mecoprop-p/A].

### WHERE TO USE:

[BONIDE Lawn Weed Killer may be used][Use] on bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, this product controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain.

### WHEN TO USE:

1. Spray during the growing season when weeds are actively growing.
2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
3. Poor weed control may result if spray is applied during drought or just before rain.
4. For best results, mow before treating.
5. Do not water lawn within 24 hours after treatment.

**HOW MUCH TO USE (Entire Lawn):****Hose End Sprayer:**

Use a hose end sprayer designed for lawns. Pour specified amount of product (Column 1) in bottle and add water to indicated level (Column 2), and spray on area shown in Column 3. For example, to spray 2,500 square feet measure 10 fluid ounces (20 tablespoons) in jar, add water to the 10 gallon level, and mix. Turn on hose and spray as evenly as possible over 2,500 square feet.

**Chart 1:** Amounts of BONIDE Lawn Weed Killer for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2 Fill Bottle to	Column 3 Area to Treat
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	5	1,250
20	10	10	2,500
30	15	15	3,750
40	20	20	5,000

Note: For the grasses in Chart 1, apply the entire contents of this container (32 fluid ounces) on 8,000 square feet of turf.

**Chart 2:** Amounts of BONIDE Lawn Weed Killer for centipedegrass and St. Augustinegrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2 Fill Bottle to	Column 3 Area to Treat
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	5	2,500
20	10	10	5,000
30	15	15	7,500
40	20	20	10,000

Note: For St. Augustinegrass and centipedegrass apply the entire contents of this container (32 fluid ounces) on 16,000 square feet of turf.

**Pressure Sprayers:**

Measure indicated amount of BONIDE Lawn Weed Killer, pour into tank, and dilute with water according to the following charts. Use a spray pattern that gives small [coarse] spray droplets without any fine mist, and cover the area indicated.

**Chart 3.** For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns with pressure or tank type sprayers.

Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer (Gallons)	Area To Treat (Square Feet)
(Tablespoons)	(Fluid Ounces)		
2	1	1	250
4	2	2	500
6	3	3	750

**Chart 4:** For centipedegrass and St. Augustinegrass lawns with pressure or tank type sprayers.

Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer (Gallons)	Area To Treat (Square Feet)
(Tablespoons)	(Fluid Ounces)		
2	1	1	500
4	2	2	1,000
6	3	3	1,500

**Spot Spraying [Treatment]:**

To spray small areas or individual weeds, do not use a hose end sprayer. Refer to Chart 3 or Chart 4 for mixing instructions. Spray as needed and be certain to wet the weeds.

<b>BROADLEAF WEEDS</b>			
Annual yellow sweetclover	Cockle	Mugwort	Speedwell
Aster	Cocklebur	Musk thistle	Spiny amaranth
Austrian fieldcress	Common mullein	Mustard	Spiny cocklebur
Bedstraw	Creeping Jenny (Creeping Charlie)	Narrowleaf plantain	Spotted catsear
Beggarticks	Cudweed	Narrowleaf vetch	Spotted spurge
Betony, Florida	Curly dock	Nettle	Spurweed
Bindweed, field	Daisy, English	Orange hawkweed	Stinging nettle
Bird vetch	Daisy, fleabane	Oriental cocklebur	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Oxalis	Tall nettle
Bittercress, hairy	Dandelion	Parsley-piert	Tall vervain
Bitterweed	Dichondra	Parsnip	Tansy ragwort
Black-eyed Susan	Dogbane	Pearlwort	Tansy mustard
Black medic	Dogfennel	Pennycress	Tanweed
Black mustard	Dollarweed	Peppergrass	Thistle
Blackseed plantain	Elderberry	Pepperweed	Trailing crownvetch
Blessed thistle	False dandelion	Pigweed	Tumble mustard
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble pigweed
Blue lettuce	False sunflower	Plains coreopsis (tickseed)	Velvetleaf
Blue vervain	Fiddleneck	Plantain	Venice mallow
Bracted plantain	Florida pusley	Poison ivy	Virginia buttonweed
Brassbuttons	Frenchweed	Poison oak	Virginia creeper
Bristly oxtongue	Galinsoga	Pokeweed	Virginia pepperweed
Broadleaf dock	Goathead	Poorjoe	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prairie sunflower	Western clematis
Broomweed	Ground ivy	Prickly lettuce	Western salsify
Buckhorn	Gumweed	Prickly sida	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate knotweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate pigweed	Wild buckwheat
Bull thistle	Healall	Prostrate spurge	Wild carrot
Bullnettle	Heartleaf drymary	Prostrate vervain	Wild four-o'clock
Burclover	Heathaster	Puncturevine	Wild garlic
Burdock	Hedge bindweed	Purslane, common	Wild geranium
Burning nettle	Hedge mustard	Ragweed	Wild lettuce
Burweed	Hemp	Red sorrel	Wild marigold
Buttercup	Henbit	Redroot pigweed	Wild mustard
Buttonweed	Hoary cress	Redstem filaree	Wild onion
Canada thistle	Hoary plantain	Rough cinquefoil	Wild parsnip
Carolina geranium	Hoary vervain	Rough fleabane	Wild radish
Carpetweed	Horsenettle	Russian pigweed	Wild rape
Catchweed bedstraw	Jimsonweed	Russian thistle	Wild strawberry
Catnip	Knawel	Scarlet pimpernel	Wild sweet potato
Catsear	Knotweed	Scotch thistle	Wild vetch
Chickweed, common	Kochia	Sheep sorrel	Woodsorrel
Chickweed, mouseear	Lambsquarters	Shepherdspurse	Woolly croton
Chicory	Lespedeza	Slender plantain	Woolly morningglory
Cinquefoil	Mallow	Smallflower galinsoga	Woolly plantain
Clover, crimson	Matchweed	Smooth dock	Wormseed
Clover, hop	Mexicanweed	Smooth pigweed	Yarrow
Clover, red	Milk vetch	Sorrel	Yellow rocket
Clover, strawberry	Morningglory	Sowthistle	Yellowflower pepperweed
Clover, sweet	Mouseear hawkweed	Spanishneedles	and other broadleaf weeds
Clover, white		Spiny sowthistle	



## **STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Keep from freezing. Store and transport in an upright position. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## **LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

For information on pesticide products (including health concerns, medical emergencies or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

### **Bonide Products, LLC**

6301 Sutliff Road  
Oriskany, NY 13424

[Lot No. added at production]

©Bonide Products, Inc. All Rights Reserved