. 4-402	D			10	16/20	03	1-	- 4	
	Environmenta	United States	-	ncy		Aregisti	ation		<u>expires 2-28-5</u> ifier Number
		Applicatio	n for I	Pesticio	le - Sectio	n I			
1. Company/Product Numbe	r Al			1 1	roduct Managa		3. P	roposed Clas	sification
	UVe	<u> </u>			a Koc	ca	—  [	None	Restricted
4. Company/Product (Name) Bonide Eight Gardi		ict Contra		PM#	13				
5. Name and Address of Applicant (Include ZIP Cade) Bonide Products, Inc. 6301 SUTTIFF RUA CL Oriskany, NY 13424 Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No Product Name						
			Sec	tion - I					
Amendment - Explain	1 below.				Final printed la Agency letter (		se to	NOTIFIC	ATION
Resubmission in resp	ionse to Agency letter below.	r dated	<u> </u>	- 🗌	"Me Too" App Other - Explain	lication.		OCT 1 (	
This neuffication is constitute with the understands that it is a violation of I 152-46, this product may be in violat	e provisions of PR Natice 98-10 : 8 U.S.C. Sec. 1001 to willfully m	ake any faise statement	10 CFR 152. 10 EPA and a enforcement	ió, land no cuhe hurther understa	r changes have been in inds that (f this notified nalities under sections	ade to the labeling o vitan is not consistent			
1. Material This Product Wil	Be Packaged In:								
Child-Resistant Packaging Yes No * Certification must be submitted	Unit Packaging Yes No If "Yes" Unit Packaging wgt	No. per . conteiner	If "Yes	Soluble Pa Yes No ;* ge wgt	No. per container	2. Type o	f Containe Matal Plastic Glass Paper Other (	r (Specify)	
3. Location of Net Contents	Information	4. Size(s) Rate	] il Contei		L	Location of L	hel Direct		
	Container					]			
6. Menner in Which Label is		Lithogr Paper of Stencil	aph lued ed		Other				
Section - IV									
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)									
Kate Entwistie Registrati			n Spec	ialst	Telephon 315-	ne No. Urotua 7 <u>36-7</u> 2	le Aree Code) 231		
Certification I certify that the statements I have made on this form and all attachme I acknowledge that any knowlingity false or misleading statement may both under applicable law.							6. Liste Ap Réceive (Sta	-	
2. Signature	<u> </u>	3	. Tido			······································		- ~ ~ ~	
Karte Fr	tuste	. IT	Realis	tratic	n Sher	valist		• • •	
4. Typed Name	······································	5	. Date				· <u>·····</u> ····	1	
Kate Entr	vislle			9-2	29-03	3			

EPA	Form	8570-1	(Rev. 3-94)	Previous	editions (	ire obsolete.

NOTIFICATION

OCT 1 6 2003



# GARDEN & HOME INSECT CONTROL

# Ready to Use

[ADDITIONAL BRAND NAME(S): Rose & Flower Ready to Use; Tomato & Vegetable Ready to Use; Home Pest Control Ready to Use; EIGHT Insect Control Garden & Home]

- READY-TO-USE
- IT'S THE LEAST YOU CAN DO FOR YOUR GARDEN.
- EASY TO USE

١

- FOR USE ON PLANTS.
- GENTLE TO MOST PLANTS.
- SAFE TO USE ON OVER 60 VARIETIES OF ORNAMENTALS\*.
- FOR CONTROL OF VEGETABLE PESTS.
- CONTROLS MOST ORNAMENTAL PESTS.
- CONTROLS MOST GARDEN INSECT PESTS.
- USE INDOORS AND OUTDOORS.
- FOR USE IN AND AROUND THE HOME..
- FOR CONTROL OF VARIOUS PLANT PESTS.
- FOR PLANT PEST CONTROL.
- WATER-BASED.
- USE AROUND BUILDINGS, HOME, FLOWER AND VEGETABLE GARDENS, AND ON FRUIT, NUT AND ORNAMENTAL TREES.
  - FOR RESIDENTIAL USE ONLY

### **KILLS OVER 100 DIFFERENT NAMED INSECTS**

\*When Used According to label instructions.

Protects your Roses, Flowers, and Vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Bagworms, Beetles, Borers, Brown dog ticks, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall Cankerworms, Fire ants, Fleas, Green fruitworms, Gnats, Grasshoppers, Houseflies, Inchworms, Japanese beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybuga, Millipedes, Mole crickets, Mosquitoes, Moths, Navel Orange worms, Scales, Sod webworms, Sowbugs (pillbugs), Spider mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies and many othera.

ACTIVE INGREDIENTS: Permethrin	
KEEP OUT OF REACH OF CHILDREN	4 • • 5 • • • • • •
CAUTION	· • 9.1
(See Side/Back Panel for Additional Precautionary Statemer Net Contents 32 FL, OZ. (1 QT.) 128 FL, OZ.	nts)
EPA Reg. No. 4-406	EPA Est. No. 4-NY-1

## PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Wash after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

### Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

#### SHAKE WELL BEFORE USING

## STORAGE AND DISPOSAL

Store and transport in an upright position. Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry area away from heat or open flame. Do not reuse container. Rinse thoroughly and wrap container in several layers of newspaper and discard in trash.

For use, turn nozzle to spray position, point away from face, and squeeze trigger to spray.

#### **INDOOR APPLICATION**

Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

#### HOUSEPLANTS

#### INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES, AND FOR INDOOR LANDSCAPING:

To kill insects on Indoor Landscaping direct the spray mist to hit as many insects as possible.

Pests:	How to Use:
Aphids, Amyworms, Citrus Blackfly, Com carworms, Exposed thrips, Fall canterworms, Inchworms, Japanese bortles, Lasthniners, Lestfollers, Mealybags, Scales, Solder mitts, Spring canterworms, Whiteflies.	Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Apply in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore a wide variety of susceptibility to injury. To ascertain possible
Ameryworms, Calibagewerns, Calibagewerns, Lastrollers	<ul> <li>phytotoxicity, spray a few plants as described above, wait</li> <li>24 hours, and check for possible injury. Repeat</li> <li>applications may be necessary to control citrus blackflies</li> <li>on non-bearing citrus, Adult Japanese beetles on roses</li> </ul>
-	Aphids, Armyworms, Citrus Blackfly, Corn carworms, Exposed thrips, Fall cashterworms, Japanese bootles, Lasfininers, Lesfollers, Mealybogs, Scales, Solider mitte, Spring cashterworms, Whiteflies. Annyworms, Cabboguwams,

Velvet Plants

Fungus gnat

EPA Reg. No. 4-406

EFA Est. No. 4-NY-1

## **INDOOR APPLICATION**

For spot application only. Do not use as a space spray indoors. Do not apply on or into electrical switch boxes, wiring or outlets. Do not allow children or pets on treated surfaces until spray has completely dried and area is ventilated. Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

[OPTIONAL DIRECTIONS FOR USE FOR REFILLABLE PACKAGING USING BONIDE" EIGHT" INSECT CONTROL CONCENTRATE, EPA REG. NO. 4-408]

**REFILLING:** This pump sprayer bottle can be refilled with BONIDE\* EIGHT\* Insect Control Concentrate. Add 1 fl. oz. of concentrate and dilute with 1 gallon of water. (Add ¼ fl. oz. of concentrate and dilute with 1 quart of water.) Apply cap tightly and *shake well to mix*. Do not use any product other than BONIDE EIGHT Insect Control Concentrate to refill this container.

USE AS A SURFACE RESIDUAL SPRAY FOR CONTROL OF: ANTS, BEDBUGS, BROWN DOG TICKS, CADELLES, CARPENTER ANTS, CARPET BEETLES, CENTIPEDES, CHEESE MITES, CLOTHES MOTHS, CLOVER MITES, COCKROACHES, CONFUSED FLOUR BEETLES, CRICKETS, DRUGSTORE BEETLES, EARWIGS, FIREBRATS, FLEAS, FLYING MOTHS, GNATS, GRAIN MITES, GRANARY WEEVILS, HOUSE FLIES, LICE AND LOUSE EGGS, LONE STAR TICKS, MEAL WORMS, MILLIPEDES, MOSOUITOES, PALMETTO BUGS, PILLBUGS, RUST RED FLOUR BEETLES, RICE WEEVILS, SCORPIONS, SILVER FISH, SOWBUGS, SPIDERS AND WATERBUGS: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. To reduce annoyance from Mosqueoes, Houseflies. Gnats and Flying Moths, apply every 2 weeks to the outside surfaces of screens, doors, window frames and outdoor tight fixtures.

TO CONTROL CARPET BEETLES: Spray mist edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed. Do not allow children or pets in treated areas until spray has dried.

<u>TO CONTROL BEDBUGS</u>: Spray mist mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed. Do not use sprayed areas until completely dry.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Do not allow children or pets to access treated surfaces until completely dry.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

EPA Reg. No. 4-406

(

EPA Est. No. 4-NY-1

## OUTDOOR APPLICATION DIRECTIONS FOR APPLYING ROSES & FLOWERS

1 1	Pests:	How to Use:	•
Ageratum, Aster, Ardisia, Azalca, Baby's breath, Begonia, Birds-nest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of thorns, cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits foot fern. Rhododendron, Rose, Schefflera, Snapdragon, Snake	Aphida, armyworms, Bagworms, Cabbage worms, Cicadas, Cornworms, cutworms, Exposed thrips, Fall cankerworms, Fall webworms, Fungus gnar, Inchworms, Japanese beetles, Leafminers, Leaffollers, Mealybugs, Scale crawlers, Spring cankerworms, Spider mites, Whiteflies, Rose Chafer, Psyllids, Scale Insects.	Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection.	
Plant, Statice, Velvet Plant, Verbena, Zinnia	5	<u> </u>	\$

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

## **VEGETABLES**

	Pests:	How to Use:
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Apply as needed to ensure uniform coverage. Can be applied up to 3 days prio- to harvest.
(posi-harvesi)	Larval stages of asparagus beetle; tamished plant bug; Lygas bugs and adult Japanese ; beetle.	Treat fems or bush growth after spear harvest when insects are present
Beans	Mexican bean beetle, Leafhoppers, Fiea beetles, Corn servorm, Lygus and stinkbugs.	Apply as needed. Can be applied up to 3 days prior to harvest.
Celery - See Leafy Vegetables		1
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworm, Cabbage sphid, Cabbage looper, Diamond back moth, imported cabbageworm.	Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest.
Com - See Sweet Com		
Cucurbits: Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds, melons including hybrids such as cantaloupe, casaba, crenshaw, honcydew, honcy balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon.	Aphids, Cabbage looper, Cucumber beetle (adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant bugs, Squash bugs.	Apply as needed to ensure uniform coverage. Can be applied up to the day of the harvest
Eggplant	Colorado potato beetles, Vegetable leafminers.	Apply as needed to ensure uniform coverage. Can be applied up to 3 1-y2 priot to harvest

EPA Reg. No. 4-406

)

EPA Est. No 4-NY-1

۰. د. د

•

4

	Pests:	How to Use:
Horseradish (Foliar application)	Imported crucifer weevils.	Apply as needed. Can be applied up to 22 days prior to harvest.
Leaft Venetables: Amaranth, Arrugula, Celery, Celtuce, Chervil, Com salad, Chrysanthemum (edible-leaved and garland), cress (garden and upland), Dandelion, Dock, Endive, Feanel, Lettuce (head and leaf), Okra, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard.	Alfalfa looper, Aphids, Boet Armyworm. Cabbage looper, Carn eer worm. Cutworms, European com borer, Fall armyworm, Green cloverworm, Southern armyworm, Tobacco budworm, Vegetable leaf miner.	Apply every five to ten days as needed. Do not make more than seven applications per crop. Can be applied up to one day prior to harvest.
Melons - See Cucurbits.		
Potatoes	Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, carworms, European com borer, Potato psyflid, Tamish plant bug.	Apply as needed. Can be applied up to 7 days prior to harvest.
Peppers	Vegetable leafminers, Cabbage looper, Flea beste, Com carworn, Pepper weevil, European com borer.	Apply as needed. Can be applied up to 3 days prior to harvest.
Strawberries, Raspberries, Blueberries.	Strawberry weevils, Aphids, Spider mites. Lygus bugs, Flea baseles, Omnivorous leaftier, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leaftoller, Weevils.	Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
Sweet Corn	Com carworm, European com borer, Fall armyworm, Southern armyworm	Apply every 5 days and when needed, out not more than 6 applications per season Can be applied up to 1 day prior to harvest
Tomatoes	Beet and southern annyworm, Tomato pinworm, Tomato fluitworm, Horn worms. Granulate cutworms, Colorado potato beetle, Cabbase looper.	Apply as needed for uniform coverage Ca be applied up to day of harvest

## VEGETABLE (CONT.)

# HOME OWNER GROWN FRUIT AND NUT TREES Apply to trees until foliage is dripping wet. Make sure to spray the undersides of leaves and into dense foliage.

Trees (fruit or aut);	Target Pests:	Use Directions:	
Almonds, Pecans	Navel orange worm: Peach twig borer.	Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.	
Apples	Plum curculio; redbanded Leafroller, Rasyappie aphid. spotted tentiform Leafminer, Tarnished plant bug. White apple Leafroller, oblique banded Leafroller. Green fruitworm.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fail	
Peaches	Lesser peach tree borer, Peach twng borer, Plum curculio, Tarnished plant bug, Gross fluit worm, Oriental fruit moth.	Apply when insects first appear Doing apply within 7 days of harvest. Do not make more than 8 applications per season.	

## EPA Reg. No. 4-406

(

(

<u>ь</u> вес:

. .

--- 6-9

Pears	Pear psylla, Green fruitworm and Codling moth.	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to
		14 days prior to harvest.
	NUTCANCE DECTS DI OUTDOOD	ADEAC

7-(1

## NUISANCE PESTS IN OUTDOOR AREAS

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

#### **OUTSIDE SURFACES OF BUILDING**

Use for control of Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

## **OTHER OUTDOOR AREAS**

(

Use for control of Ants, Crickets, Earwigs, Fleas, Lone star ticks for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

#### Fire Ant Control Individual Mound and Plant Container Treatment

Gently apply over the surface of each mound. Thoroughly wet mound and surrounding area of 3 x 4 ft. (12 sq. ft.) For best results apply in cool weather 65°F-80°F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration, which will reduce product effectiveness. Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.

For use only as an aid in reducing annoyance from these insects on the following: Cluster flies, Flying moths, Gnats, House flies, Mosquitoes. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

For information on pesticide products (including health concerns, methoal emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by Bonide Products, Inc. Oriskany, NY 13424

EFA Est. No. 4-NN-1

EPA Reg. No. 4-406

## Optional Marketing Statements (Continued from page 1) EPA Reg. No. 4-406

8-9

Protects your Houseplants, Roses, Flowers, Vegetables, Fruit, and Nuts from: Alfalfa Loopers, Ants, Aphids, Armyworms, Asian Ladybugs, Bagworms, Beetles, Borers, Budworms, Cabbageworms, Chinch Bugs, Cicadas, Citrus Black Flies, Codling Moths, Corn Earworms, Cutworms, Elm Spanworms, Fall Cankerworms, Green Fruitworms, Gnats, Grasshoppers, Inchworms, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Lygus bugs, Mealybugs, Mole Crickets, Navel Orange Worms, Scales, Sod Webworms, Sowbugs (pillbugs), Spider Mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies, and many others.

Protects your home from: Carpenter Ants, Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clovermites, Cheese Mites, Granary Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites, Cadelles, Fire Ants, Crickets, Earwigs, Fleas, and Lone Star Ticks.

Use in and around the home on...Houseplants, Vegetables, Roses, Flowers, Ornamentals

# **Bonide Products, Inc.**

6301 Sutliff Road Orlskany, NY 13424

Telephone: 315-736-8231 Fax: 315-736-7582 Email: mattv@bonide.com

September 29, 2003

1

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) US Environmental Protection Agency Room 266A, Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202-4501

## RE: Bonide Eight Garden & Home Insect Control EPA Reg. 4-406 Notification of optional marketing statements:

Protects your Houseplants, Roses, Flowers, Vegetables, Fruit, and Nuts from: Alfalfa Loopers, Ants, Aphids, Armyworms, Asian Ladybugs, Bagworms, Beetles, Borers, Budworms, Cabbageworms, Chinch Bugs, Cicadas, Citrus Black Flies, Codling Moths, Corn Earworms, Cutworms, Elm Spanworms, Fall Cankerworms, Green Fruitworms, Gnats, Grasshoppers, Inchworms, Japanese Beetles, Leafnoppers, Leafnoilers, Lygus bugs, Mealybugs, Mole Crickets, Navel Orange Worms, Scales, Sod Webworms, Sowbugs (pillbugs), Spider Mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies, and many others.

Protects your home from: Carpenter Ants, Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clovermites, Cheese Mites, Granary Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites, Cadelles, Fire Ants, Crickets, Earwigs, Fleas, and Lone Star Ticks.

Use in and around the home on ... Houseplants, Vegetables, Roses, Flowers, Omamentals

To Whom It May Concern:

Enclosed, please find a completed form 8570-1, Notification for optional marketing statements and labeling for the above-mentioned registration. I trust you will find everything in order.

Should you have any questions or need additional information, please do not hesitate to contact me at (315) 736-8231 ext. 230.

<u>د</u> به

Sincerely,

anthtusen

Kate Entwistle Registration Specialist