11 1100

2006/06/50

1/10

,

4-406		Ujac	12000			10
Please read instructions on	reverse before completing	form.	Form Appro	ved. OMB No. 20	70-006	D. Approvel expires 2-28-9
\$-EPA	Environmental P	od States rotection Agenc on, DC 20460	y [Registrat Amendm Other	ion	OPP Identifier Number
	Ap	plication for Pe	sticide - Section	on I		<u> </u>
1. Company/Product Number 4-406		2.	EPA Product Manag aRocca		3. Pro	oposed Classification
4. Company/Product (Name Bonide Eight Garden &			A# 3			None Restricted
5. Name and Address of Ap	plicant (Include ZIP Code)	6.	Expedited Revei	w. In accordance	ce with	FIFRA Section 3(c)(3)
Bonide Products, Inc 6301 Sutliff Road Oriskany, NY 13424		(b to)(i), my product is	similar or identic	al in coi	mposition and labeling
Check if thi	s is a new address	P	roduct Name			
	······································	Sectio	n - II			
Amendment - Explai	n below. bonse to Agency letter dat	ed	Final printed I Agency latter "Me Too" Ap	-		OTIFICATION
Notification - Explain	below.		Other - Explai	n below.	M	IAR 2 0 2006
and Bonide Products, Inc. m		nt action and penalties un Sectior		ot FIFRA.		
1. Material This Product Wil	·			·····		
Child-Resistant Packaging	Unit Packaging		ible Packaging	2. Type of C	entainer Metal	
Yes No	Yes No	Yes No			Plastic	
* Certification must be submitted	lf "Yes" N	o. per If "Yes ontainer Package w	gt Container		Glass Paper Other (S	pecify)
3. Location of Net Contents	Information 4. 1	Size(s) Retail Container	5	Location of Label	Directio	ns
6. Manner in Which Label is	Affixed to Product		Other	· · · · · · · · · · · · · · · · · · ·		
		Section	- IV			
1. Contact Point (Complete	items directly below for id	lentification of individua	to be contacted, if i	necessary, to proc	ess this	application.)
Name Nicole Russell		Title Registration	Specialist		•	No. (Include Area Code) 231 ext 230
	ments I have made on this y knowlingity false or misi					6. Date Application Received (Stamped)
2. Signature Nigot	ussell	7 3. Title Registration S	Specialist			
I. Typed Name		5. Date				
Nicole Russell	/	ļ	03-10-06			

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.



GARDEN & HOME INSECT CONTROL

Ready to Use

[ADDITIONAL BRAND NAME(S): Rose & Flower Ready to Use; Tomato & Vegetable Ready to Use; Home Pest Control Ready to Use; EIGHT Insect Control Garden & Home]

·READY-TO-USE ·IT'S THE LEAST YOU CAN DO FOR YOUR GARDEN. -EASY TO USE FOR USE ON PLANTS. -GENTLE TO MOST PLANTS. -SAFE TO USE ON OVER 60 VARIETIES OF ORNAMENTALS*. ·FOR CONTROL OF VEGETABLE PESTS. ·CONTROLS MOST ORNAMENTAL PESTS. -CONTROLS MOST GARDEN INSECT PESTS. **USE INDOORS AND OUTDOORS.** FOR USE IN AND AROUND THE HOME ... ·FOR CONTROL OF VARIOUS PLANT PESTS. FOR PLANT PEST CONTROL. ·WATER-BASED. -USE AROUND BUILDINGS, HOME, FLOWER AND VEGETABLE GARDENS, AND ON FRUIT, NUT AND ORNAMENTAL TREES. -FOR RESIDENTIAL USE ONLY **KILLS OVER 100 DIFFERENT NAMED INSECTS**

*When Used According to label instructions.

Protects your Roses, Flowers, and Vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Asian ladybug, Bagworms, Beetles, Borers, Brown dog ticks, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths. Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall Cankerworms, Fire ants, Fleas, Green fruitworms, Gnats. Grasshoppers, Houseflies, Inchworms, Japanese beetles. Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Moths, Navel Orange worms. Scales, Sod webworms, Sowbugs (pillbugs), Spider mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies and many others.

ACTIVE INGREDIENTS:

Permethrin	(Cas. No. 52645-53-1)	0.02%
OTHER INGREDIENTS:		<u>99.98%</u>
TOTAL		

KEEP OUT OF REACH OF CHILDREN CAUTION

(See Side/Back Panel for Additional Precautionary Statements) Net Contents 32 FL. OZ. (1 QT.) 128 FL. OZ. (1 GAL.) PRECAUTIONARY STATEMENTS

EPA Reg. No. 4-406 3/10/06 - notification EPA Est. No. 4-NY-1 Page 1 of 7 Pages Hazards to Humans and Domestic Animals

CAUTION: Wash after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry area away from heat or open flame.

Disposal: If empty - Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For use, turn nozzle to spray position, point away from face, and squeeze trigger to spray.

INDOOR APPLICATION

Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

HOUSEPLANTS INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES, AND FOR INDOOR LANDSCAPING:

To kill insects on Indoor Landscaping direct the spray mist to hit as many insects as possible.

Plants:	Pests:	How to Use:
Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Citrus, Coleus, Crown of Thorns. Cockscornb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Citrus Blackfly, Corn earworms, Exposed thrips, Fall cankerworms, Japanese beetles. Leafminers, Leafrollers, Mealybugs, Scales, Solder mites, Spring cankerworms. Whiteflies.	Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Apply in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemurns, and roses, there exists a wide variety of species and cultivars, and therefore a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury. Repeat applications may be necessary to control citrus blackflies on non-bearing citrus. Adult Japanese beetles on roses.
Chrysanthemums	Armyworms, Cabbageworms, Cutworms, Leafrollers	
Velvet Plants	Fungus gnats]

INDOOR APPLICATION

EPA Reg. No. 4-406 3/10/06 - notification EPA Est. No. 4-NY-1 Page 2 of 7 Pages **For spot application only.** Do not use as a space spray Indoors. Do not apply on or into electrical switch boxes, wiring or outlets. Do not allow children or pets on treated surfaces until spray has completely dried and area is ventilated. Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

[OPTIONAL DIRECTIONS FOR USE FOR REFILLABLE PACKAGING USING BONIDE" EIGHT" INSECT CONTROL CONCENTRATE, EPA REG. NO. 4-408]

REFILLING: This pump sprayer bottle can be refilled with BONIDE⁶⁰ EIGHT⁶⁰ Insect Control Concentrate. Add 1 fl. oz. of concentrate and dilute with 1 gallon of water. (Add ¹/₄ fl. oz. of concentrate and dilute with 1 quart of water.) Apply cap tightly and *shake well to mix*. Do not use any product other than BONIDE EIGHT Insect Control Concentrate to refill this container.

USE AS A SURFACE RESIDUAL SPRAY FOR CONTROL OF: ANTS, ASIAN LADYBUGS, BEDBUGS, BROWN DOG TICKS, CADELLES, CARPENTER ANTS, CARPET BEETLES, CENTIPEDES, CHEESE MITES, CLOTHES MOTHS, CLOVER MITES, COCKROACHES, CONFUSED FLOUR BEETLES, CRICKETS, DRUGSTORE BEETLES, EARWIGS, FIREBRATS, FLEAS, FLYING MOTHS, GNATS, GRAIN MITES, GRANARY WEEVILS, HOUSE FLIES, LICE AND LOUSE EGGS, LONE STAR TICKS, MEAL WORMS, MILLIPEDES, MOSQUITOES, PALMETTO BUGS, PILLBUGS, RUST RED FLOUR BEETLES, RICE WEEVILS, SCORPIONS, SILVERFISH, SOWBUGS, SPIDERS AND WATERBUGS: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. To reduce annoyance from Mosquitoes, Houseflies, Gnats and Flying Moths, apply every 2 weeks to the outside surfaces of screens, doors, window frames and outdoor light fixtures.

<u>TO CONTROL CARPET BEETLES</u>: Spray mist edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed. Do not allow children or pets in treated areas until spray has dried.

TO CONTROL BEDBUGS: Spray mist mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed. Do not use sprayed areas until completely dry.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Do not allow children or pets to access treated surfaces until completely dry.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

OUTDOOR APPLICATION

EPA Reg. No. 4-406 3/10/06 - notification EPA Est. No. 4-NY-1 Page 3 of 7 Pages

DIRECTIONS FOR APPLYING ROSES & FLOWERS

	Pests:	How to Use:
Ageratum, Aster, Ardisia, Azalea, Baby's	Aphids, armyworms,	Apply as a thorough spray, wetting
breath, Begonia, Birds-nest fern, Bleeding	Bagworms, Cabbage	the plants to the dripping point. Try
heart, Carnation, Chrysanthemum, Coleus,	worms, Cicadas,	to hit underside of leaves and
Common Ninebark and Snowberry, Crown of	Cornworms, cutworms,	penetrate dense foliage. Spray in
thorns, cockscomb, Cyclamen, Dracaena,	Exposed thrips, Fall	the late afternoon or evening, when
Dumbcane, English Ivy, Exacum, Fuchsia,	cankerworms, Fall	the temperature ranges from 50° to
Gladiolus, Grape Ivy, Gold Bells, Hypoestes,	webworms, Fungus gnat,	75°F and when there is little or no
Ivy, Juniper, Lilac, Marigold, Mock-Orange,	Inchworms, Japanese	wind. Spray at the first sign of
Nannyberry, Orchid, Palm, Pansy, Pea Shrub,	beetles, Leafminers,	insects. Repeat as necessary; use
Peperomia, Petunia, Philodendron, Piggy-back	Leafrollers, Mealybugs,	intervals of 4-8 days. Application
Plant, Poinsettia, Portulaca, Pothos, Prayer	Scale crawlers, Spring	can be made up to the day of
Plant, Purple Passion, Rabbits foot fern,	cankerworms, Spider	collection.
Rhododendron, Rose, Schefflera, Snapdragon,	mites, Whiteflies, Rose	
Snake Plant, Statice, Velvet Plant, Verbena,	Chafer, Psyllids, Scale	
Zinnia	Insects.	

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

	Pests:	How to Use:
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Apply as needed to ensure uniform coverage. Can be applied up to 3 days prior to harvest.
(post-harvest)	Larval stages of asparagus beetle; tarnished plant bug; Lygus bugs and adult Japanese beetle.	Treat ferns or bush growth after spear harvest when insects are present.
Celery - See Leafy Vegetables	¥	Y
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworm, Cabbage aphid. Cabbage looper, Diamond back moth, Imported cabbageworm.	Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest.
Corn - See Sweet Corn		
Cucurbits: Balsam pear (bitter melon). Chinese waxgourd (Chinese preserving melon), citron melon. cucumber, gherkin, edible gourds, melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon.	Aphids, Cabbage looper, Cucumber beetle (adults), Cutworms, Leafhoppers, Leafminers. Melonworms, Pickleworm, Plant bugs, Squash bugs.	Apply as needed to ensure uniform coverage. Can be applied up to the day of the harvest.
Eggplant	Colorado potato beetles, Vegetable leafminers.	Apply as needed to ensure uniform coverage. Can be applied up to 3 days prior to harvest.

VEGETABLES

VEGETABLE (CONT.)

	Pests:	How to Use:
Horseradish	Imported crucifer weevils,	Apply as needed. Can be applied

EPA Reg. No. 4-406 3/10/06 - notification

I

EPA Est. No. 4-NY-1 Page 4 of 7 Pages

Deleted: Beans

Deleted: Mexican bean beetle, Leafboppers, Flea beetles, Corn earworm, Lygus and stinkbugs.

Deleted: Apply as needed. Can be applied up to 3 days prior to harvest.

(Foliar application)	·····	up to 22 days prior to harvest.
Leafy Vegetables: Amaranth, Arrugula, Celery, Celtuce, Chervil, Com salad, Chrysanthemum (edible-leaved and garland), cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Okra, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard.	Alfalfa looper, Aphids, Beet Armyworm, Cabbage looper, Corn ear worm, Cutworms, European corn borer, Fall armyworm, Green cloverworm, Southern armyworm. Tobacco budworm, Vegetable leaf miner.	Apply every five to ten days as needed. Do not make more than seven applications per crop. Can be applied up to one day prior to harvest.
Melons - See Cucurbits.		
Potatoes	Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, cutworms, European corn borer, Potato psyllid, Tarnish plant bug.	Apply as needed. Can be applied up to 7 days prior to harvest.
Peppers	Vegetable leafminers, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer.	Apply as needed. Can be applied up to 3 days prior to harvest.
Strawberries, Raspberries, Blueberries.	Strawberry weevils, Aphids, Spider mites, Lygus bugs, Flea beetles, Omnivorous leaftier, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leafroller, Weevils.	Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
Sweet Corn	Corn earworm, European corn borer, Fall armyworm, Southern armyworm.	Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomatoes	Beet and southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle, Cabbage looper.	Apply as needed for uniform coverage. Can be applied up to day of harvest.

HOME OWNER GROWN FRUIT AND NUT TREES

Apply to trees until foliage is dripping wet. Make sure to spray the undersides of leaves and into dense foliage.				
Trees (fruit or nut):	Target Pests:		Use Directions:	

EPA Reg. No. 4-406 3/10/06 - notification EPA Est. No. 4-NY-1 Page 5 of 7 Pages

Almonds, Pecans	Navel orange worm; Peach twig borer.	Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Plum curculio; redbanded Leafroller, Rosyapple aphid, spotted tentiform Leafminer, Tarnished plant bug, White apple Leafroller, oblique banded Leafroller, Green fruitworm.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
Peaches	Lesser peach tree borer, Peach twig borer. Plum curculio, Tarnished plant bug, Green fruit worm, Oriental fruit moth.	Apply when insects first appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears	Pear psylla, Green fruitworm and Codling moth.	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.

NUISANCE PESTS IN OUTDOOR AREAS

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

OUTSIDE SURFACES OF BUILDING

Use for control of Ants, Asian ladybugs, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

OTHER OUTDOOR AREAS

Use for control of Ants, Crickets, Earwigs, Fleas, Lone star ticks for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

<u>Fire Ant Control</u> <u>Individual Mound and Plant Container</u> <u>Treatment</u>

Gently apply over the surface of each mound. Thoroughly wet mound and surrounding area of 3 x 4 ft. (12 sq. ft.) For best results apply in cool weather 65°F-80°F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration, which will reduce product effectiveness. <u>Use the same</u> method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.

For use only as an aid in reducing annoyance from these insects on the following: Cluster flies, Flying moths, Gnats, House flies, Mosquitoes. Spray outside surfaces of screens, doors, window frames or wherever these insects may

EPA Reg. No. 4-406 3/10/06 - notification EPA Est. No. 4-NY-1 Page 6 of 7 Pages enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by Bonide Products, Inc. Oriskany, NY 13424

Optional Marketing Statements (Continued from page 1)

Protects you Houseplants, Roses, Flowers, Vegetables, Fruit, and Nuts from: Alfalfa Loopers, Ants, Aphids, Armyworms, Asian Ladybugs, Bagworms, Beetles, Borers, Budworms, Cabbageworms, Chinch Bugs, Cicadas, Citrus Black Flies, Codling Moths, Corn Earworms, Cutworms, Elm Spanworms, Fall Cankerworms, Green Fruitworms, Gnats, Grasshoppers, Inchworms, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Lygus bugs, Mealybugs, Mole Crickets, Navel Orange Worms, Scales, Sod Webworms, Sowbugs (pillbugs), Spider Mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies, and many others.

Protects your home from: Carpenter Ants, Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clovermites, Cheese Mites, Granary Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites, Cadelles, Fire Ants, Crickets, Earwigs, Fleas, and Lone Star Ticks.

Use in and around the home on...Houseplants, Vegetables, Roses, Flowers, Ornamentals

EPA Reg. No. 4-406 3/10/06<u>- notification</u> EPA Est. No. 4-NY-1 Page 7 of 7 Pages

Bonide Products, Inc.

6301 Sutliff Rd. Oriskany, NY 13424

Phone (315) 736-8231 Fax (315) 736-7582

March 10, 2006

FedEx: 8535 5225 1014

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) US Environmental Protection Agency Room 266A, Crystal Mall 2 1801 South Bell Street Arlington, VA 22202-4501

RE: Bonide Eight Garden & Home Insect Control EPA Reg. No. 4-406 Notification – To remove beans from the Directions for Use site, per EPA's letter dated March 2, 2006.

To Whom It May Concern:

Enclosed, please find a completed form 8570-1, Notification and the latest final printed label for the above-mentioned registration. Changes are highlighted in yellow. I trust you will find everything in order for approval.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Should you have any questions or require additional information, please do not hesitate to contact me at (315) 736-8231, extension 230 or by e-mail at <u>Nicoler@bonide.com</u>.

Sincerely

Nicole Russell Registration Specialist

Enclosure





UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460



MAP 0 6 2006

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAR 2 2006

Ms. Kate Entwistee Bonide Products 6301 Sutliff Road Oriskany, NY 13424

Dear Ms. Entwistee:

Subject: Label change to remove beans Bonide Eight Garden & Home Insect Control EPA Registration No. 4-406

It has come to the Agency's attention that there is no tolerance for permethrin on beans. In the Agency last accepted label dated October 3, 2001 for your product beans were approved as a site.

Please remove beans from the label and send the Agency a copy of the new label.

Sincerely yours,

Led A Que Leven

George T. LaRocca Product Manager (13) Insecticide Branch Registration Division (7505)