1/9

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

0CT - 62009

Bonide Products, Inc. c/o Ms. Sarah Kuetemeier Registrations by Design, Inc. P.O. Box 1019 Salem, VA 24153

Dear Ms. Kuetemeier:

Subject: Application for Pesticide Notification (PRN 98-10)

Comply with Agency pyrethroid labeling letter dated June 4, 2009

Bonide Eight Insect Control Garden & Home Ready to Use

EPA Reg. No. 4-406

Your submission dated August 20, 2009

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated August 20, 2009 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Sincerely,

Dara R. Pelitt

Richard Gebken Product Manager 10

Insecticide Branch

Registration Division (7505P)

Jase read instructions on reverse before con ng form.	Form Assess	OMB No. 2070 008	49
United States Environmental Protect Washington, DC 2	ion Agency	Registration Amendment	O. Approvel expires 2-28-95 OPP Identifier Number
Applicat	ion for Pesticide - Section	n I	
1. Company/Product Number 4-406	2. EPA Product Manage Gebken	3. Pr	oposed Classification
Company/Product (Name) Bonide® Eight Insect Control Garden & Home Insect Control Re	PM# eady to Use 13		
5. Name and Address of Applicant (Include ZIP Code) Bonide Products, Inc. 6301 Sutliff Road Oriskany, NY 13424 Check if this is a new address	(b)(i), my product is so to:	v. In accordance with imilar or identical in co	mposition and labeling
	Section - II		
Amendment - Explain below. Resubmission in response to Agency letter dated The Too" Application. Notification - Explain below. Capplanation: Use additional page(s) if necessary. (For section I and Section II.)			
NOTIFICATION: of label change per June 4, 2009 letter from EPA This notification is consistent with the guidance in the June 4, 2009 letter and the requirements of EPA's regulations at 40 CFR part 156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. Bonide Products, Inc. understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA, and further understands that if the amended label is not consistent with the requirements of 40 CFR part 156, this product may be in violation of FIFRA and Bonide Products, Inc. may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.			
4 San Carlot Barrier B	Section - III		
1. Material This Product Will Be Packaged In: Child-Resistant Packaging Yes No ** Certification must be submitted Vill Packaging Unit Packaging Vinit Packaging wgt. No. per Unit Packaging wgt.	Water Soluble Packaging Yes No If "Yes" Package wgt No. per Container	2. Type of Container Metal Plastic Glass Paper Other (S	
3. Location of Net Contents Information 4. Size(s) R Label Container	etail Container 5. QT; GAL	Location of Label Direction	ons
6. Manner in Which Lebel is Affixed to Product Lithe Paper Ster	ograph Other er glued aciled		0000
Section - IV			
Contact Point (Complete items directly below for identification) Name Sarah H. Kuetemeier	Title Consultant/Agent for Bonide Products, Inc.		e No. (Include Area Code)
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowlingly false or misleading statement may be punishable by fine or imprisonment or complete. both under applicable law. 6. Date Application Received (Stamped)			
2. Signature Darah W. Kuetemerei	3. Title Consultant/Agent for Bonide Produc	sts, Inc.	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
4. Typed Name Sarah H. Kuetemeier	5. Date Aug. 20, 2009		



118½ East Main Street Suite 1 Salern, VA 24153-3805 phone 540.375.8826 fax 540.375.8827 Aug. 20, 2009

Document Processing Desk (NOTIF-PYRETHROID)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Ariel Rios Bldg.
1200 Pennsylvania Ave., NW
Washington, DC 20460

Re: Notification of Revised Label per EPA 6-4-09 Letter

EPA Reg. No. 4-406, Bonide® Eight Insect Control Garden & Home Insect Control Ready to Use

Attn: Linda Arrington, Team Leader

Notifications & Minor Formulations

On behalf of Bonide Products, Inc. of Oriskany, New York, enclosed is notification for the above-referenced product. The purpose of this notification is to update the label per the June 4, 2009 letter from EPA. No changes have been made to the CSF.

The following documents are enclosed:

- 1) EPA Form #8570-1, Application for Pesticide Notification (2 copies)
- 2) Copy of the proposed revised label with revisions highlighted in blue.

If further action or information is required on behalf of Bonide Products, Inc., please advise.

Best regards,

Sarah H. Kuetemeier

Consultant/Agent for

BONIDE PRODUCTS, INC.

buch H. Kuetemerer

e-mail: sarahk@ntelos.net

/shk

Enclosures

cc: MVickers/BPI

NOTICE! MAILING ADDRESS CHANGE effective August 1, 2009

Registrations by Design, Inc., P.O. Box 1019, Salem, VA 24153

Established 1996



NOTIFICATION
DCT - 6 2009



GARDEN & HOME INSECT CONTROL

Ready to Use

[ADDITIONAL BRAND NAME(S): Rose & Flower Ready to Use; Tomato & Vegetable Ready to Use; Home Pest Control Ready to Use; EIGHT Insect Control Garden & Home]

- ·READY-TO-USE
- ·IT'S THE LEAST YOU CAN DO FOR YOUR GARDEN.
- ·EASY TO USE
- ·FOR USE ON PLANTS.
- ·GENTLE TO MOST PLANTS.
- ·SAFE TO USE ON OVER 60 VARIETIES OF ORNAMENTALS*.
- ·FOR CONTROL OF VEGETABLE PESTS.
- ·CONTROLS MOST ORNAMENTAL PESTS.
- ·CONTROLS MOST GARDEN INSECT PESTS.
- ·USE INDOORS AND OUTDOORS.
- ·FOR USE IN AND AROUND THE HOME..
- ·FOR CONTROL OF VARIOUS PLANT PESTS.
- ·FOR PLANT PEST CONTROL.
- ·WATER-BASED.
- ·USE AROUND BUILDINGS, HOME, FLOWER AND VEGETABLE GARDENS, AND ON FRUIT, NUT AND ORNAMENTAL TREES.
- ·FOR RESIDENTIAL USE ONLY

KILLS OVER 100 DIFFERENT NAMED INSECTS

Protects your Roses, Flowers, and Vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Asian ladybug, Bagworms, Beetles, Borers, Brown dog ticks, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall Cankerworms, Fire ants, Fleas, Green fruitworms, Gnats, Grasshoppers, Houseflies, Inchworms, Japanese beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Moths, Navel Orange worms, Scales, Sod webworms, Sowbugs (pillbugs), Spider mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies and many others.

ACTIVE INGREDIENTS:

Permethrin(Cas. No. 52645-53	-1)
OTHER INGREDIENTS:	<u>99.98%</u>
TOTAL	100.00%
	0000 c c

KEEP OUT OF REACH OF CHILDREN CAUTION

(See Side/Back Panel for Additional Precautionary Statements and First Aid)

Net Contents 32 FL. OZ. (1 QT.) 128 FL. OZ. (1 GAI

EPA Reg. No. 4-406 8/20/09 – Notification EPA Est. No. 4-NY-1 Page 1 of 6 Pages

000000

^{*}When Used According to label instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Anima

CAUTION: Wash after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

Environmental Hazards

This pesticide is extremely toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Avoid contamination of ornamental fish ponds.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry area away from heat or open flame.

<u>Disposal:</u> If empty – Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For use, turn nozzle to spray position, point away from face, and squeeze trigger to spray.

INDOOR APPLICATION

Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

HOUSEPLANTS

INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES, AND FOR INDOOR LANDSCAPING:

To kill insects on Indoor Landscaping direct the spray mist to hit as many insects as possible.

Plants:	Pests:	How to Use:
Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Citrus, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Citrus Blackfly, Corn earworms, Exposed thrips, Fall cankerworms, Fungus gnats, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Solder mites, Spring cankerworms, Whiteflies.	Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Apply in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury. Repeat applications may be necessary to control citrus blackflies on non-bearing citrus, Adult of Japanese beetles on roses.
Chrysanthemums	Armyworms, Cabbageworms, Cutworms, Leafrollers	0000 000 000 000

INDOOR APPLICATION

For spot application only. D t use as a space spray Indoors. Do not apply r into electrical switch boxes, wiring or outlets. Do not allow children or pets on treated surfaces until spray has completely dried and area is ventilated. Do not use in commercial food handling establishments, restaurants or other areas where food is commercially prepared or processed. In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

[OPTIONAL DIRECTIONS FOR USE FOR REFILLABLE PACKAGING USING BONIDE® EIGHT® INSECT CONTROL CONCENTRATE, EPA REG. NO. 4-408]

REFILLING: This pump sprayer bottle can be refilled with BONIDE® EIGHT® Insect Control Concentrate. Add 1 fl. oz. of concentrate and dilute with 1 gallon of water. (Add ¼ fl. oz. of concentrate and dilute with 1 quart of water.) Apply cap tightly and shake well to mix. Do not use any product other than BONIDE EIGHT Insect Control Concentrate to refill this container.

USE AS A SURFACE RESIDUAL SPRAY FOR CONTROL OF: ANTS, ASIAN LADYBUGS, BEDBUGS, BROWN DOG TICKS, CADELLES, CARPENTER ANTS, CARPET BEETLES, CENTIPEDES, CHEESE MITES, CLOTHES MOTHS, CLOVER MITES, COCKROACHES, CONFUSED FLOUR BEETLES, CRICKETS, DRUGSTORE BEETLES, EARWIGS, FIREBRATS, FLEAS, FLYING MOTHS, GNATS, GRAIN MITES, GRANARY WEEVILS, HOUSE FLIES, LICE AND LOUSE EGGS, LONE STAR TICKS, MEAL WORMS, MILLIPEDES, MOSQUITOES, PALMETTO BUGS, PILLBUGS, RUST RED FLOUR BEETLES, RICE WEEVILS, SCORPIONS, SILVERFISH, SOWBUGS, SPIDERS AND WATERBUGS: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. To reduce annoyance from Mosquitoes, Houseflies, Gnats and Flying Moths, apply every 2 weeks to the outside surfaces of screens, doors, window frames and outdoor light fixtures.

TO CONTROL CARPET BEETLES: Spray mist edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed. Do not allow children or pets in treated areas until spray has dried.

TO CONTROL BEDBUGS: Spray mist mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed. Do not use sprayed areas until completely dry.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Do not allow children or pets to access treated surfaces until completely dry.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use. 6 60 000000

OUTDOOR APPLICATION DIRECTIONS FOR APPLYING

Do not water the treated area to the point of run-off. Do not make applications during rain. All coutdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- (3) Applications to building foundations, up to a maximum height of 3 feet.

EPA Reg. No. 4-406 8/20/09 – Notification EPA Est. No. 4-NY-1 Page 3 of 6 Pages

0000

•Other than applications to build foundations, all outdoor applications to impervious surfaces such as sidewalks, a driveways, patios, porches and ctural surfaces (such as windows, doors, and s) are limited to spot and crack-and-crevice applications only.

ROSES & FLOWERS

	Pests:	How to Use:
Ageratum, Aster, Ardisia, Azalea, Baby's	Aphids, armyworms,	Apply as a thorough spray,
breath, Begonia, Birds-nest fern, Bleeding heart,	Bagworms, Cabbage worms,	wetting the plants to the
Carnation, Chrysanthemum, Coleus, Common	Cicadas, Cornworms,	dripping point. Try to hit
Ninebark and Snowberry, Crown of thorns,	cutworms, Exposed thrips,	underside of leaves and
cockscomb, Cyclamen, Dracaena, Dumbcane,	Fall cankerworms, Fall	penetrate dense foliage. Spray
English Ivy, Exacum, Fuchsia, Gladiolus, Grape	webworms, Fungus gnat,	in the late afternoon or
Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac,	Inchworms, Japanese	evening, when the temperature
Marigold, Mock-Orange, Nannyberry, Orchid,	beetles, Leafminers,	ranges from 50° to 75°F and
Palm, Pansy, Pea Shrub, Peperomia, Petunia,	Leafrollers, Mealybugs,	when there is little or no wind.
Philodendron, Piggy-back Plant, Poinsettia,	Scale crawlers, Spring	Spray at the first sign of
Portulaca, Pothos, Prayer Plant, Purple Passion,	cankerworms, Spider mites,	insects. Repeat as necessary;
Rabbits foot fern, Rhododendron, Rose,	Whiteflies, Rose Chafer,	use intervals of 4-8 days.
Schefflera, Snapdragon, Snake Plant, Statice,	Psyllids, Scale Insects.	Application can be made up to
Velvet Plant, Verbena, Zinnia		the day of collection.
On Tout extermillors, and institute should be used when externillors are altered. First actived France		

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

VEGETABLES

	VEGETABLES				
	Pests:	How to Use:			
Asparagus	Asparagus beetles, Cutworms	Apply as needed to ensure uniform			
(pre-harvest)		coverage. Can be applied up to 3			
		days prior to harvest.			
(post-harvest)	Larval stages of asparagus beetle;	Treat ferns or bush growth after			
	tarnished plant bug; Lygus bugs	spear harvest when insects are			
	and adult Japanese beetle.	present.			
Celery - See Leafy Vegetables					
Cole Crops:	Beet armyworm, Cabbage aphid,	Apply every 5 days or as needed.			
Broccoli, Brussels Sprouts, Cabbage,	Cabbage looper, Diamond back	Can be applied up to 1 day prior to			
Cauliflower	moth, Imported cabbageworm.	harvest.			
Corn - See Sweet Corn					
Cucurbits:	Aphids, Cabbage looper,	Apply as needed to ensure uniform			
Balsam pear (bitter melon), Chinese	Cucumber beetle (adults),	coverage. Can be applied up to the			
waxgourd (Chinese preserving	Cutworms, Leafhoppers,	day of the harvest.			
melon), citron melon, cucumber,	Leafminers, Melonworms,				
gherkin, edible gourds, melons	Pickleworm, Plant bugs, Squash				
including hybrids such as cantaloupe,	bugs.				
casaba, crenshaw, honeydew, honey					
balls, mango melon, muskmelon,					
Persian melon, pumpkin, summer &					
winter squash, watermelon.					
Eggplant	Colorado potato beetles,	Apply as needed to ensure uniform			
	Vegetable leafminers.	coverage. Can be applied up to 3.			
		days prior to harvest.			
Horseradish	Imported crucifer weevils.	Apply as needed. Can be applied			
(Foliar application)		up to 22 days prior to harvest. o			
Leafy Vegetables:	Alfalfa looper, Aphids, Beet	Apply every live to ten days as o			
Amaranth, Arrugula, Celery, Celtuce,	Armyworm, Cabbage looper, Corn	needed. Do not make more than			
Chervil, Com salad, Chrysanthemum	ear worm, Cutworms, European	seven applications per crop. Can be			
(edible-leaved and garland), cress	corn borer, Fall armyworm, Green	applied up to one day prior to			
(garden and upland), Dandelion,	cloverworm, Southern armyworm,	harvest.			
Dock, Endive, Fennel, Lettuce (head	Tobacco budworm, Vegetable leaf	000C C C			
and leaf), Okra, Parsley, Purslane	miner.				
(garden and winter), Rhubarb,		C C C C C C C C C C C C C C C C C C C			
Spinach, Swiss chard.					

EPA Reg. No. 4-406 8/20/09 – Notification EPA Est. No. 4-NY-1 Page 4 of 6 Pages

VEGETABLE (CONT.)		
	Pests:	how to Use:
Melons - See Cucurbits.		
Potatoes	Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, cutworms, European corn borer, Potato psyllid, Tarnish plant bug.	Apply as needed. Can be applied up to 7 days prior to harvest.
Peppers	Vegetable leafminers, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer.	Apply as needed. Can be applied up to 3 days prior to harvest.
Strawberries, Raspberries, Blueberries.	Strawberry weevils, Aphids, Spider mites, Lygus bugs, Flea beetles, Omnivorous leaftier, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leafroller, Weevils.	Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
Sweet Corn	Corn earworm, European corn borer, Fall armyworm, Southern armyworm.	Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomatoes	Beet and southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle,	Apply as needed for uniform coverage. Can be applied up to day of harvest.

HOME OWNER GROWN FRUIT AND NUT TREES

Apply to trees until foliage is dripping wet. Make sure to spray the undersides of leaves and into dense foliage.

Cabbage looper.

Trees (fruit or nut):	Target Pests:	Use Directions:
Almonds, Pecans	Navel orange worm; Peach twig borer.	Apply when insects appear. Do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Plum curculio; redbanded Leafroller, Rosyapple aphid, spotted tentiform Leafminer, Tarnished plant bug, White apple Leafroller, oblique banded Leafroller, Green fruitworm.	Apply when insects appear. Repeat as required to maintain control. Do not make more than 3 applications per season. Do not apply after petal fall.
Peaches	Lesser peach tree borer, Peach twig borer, Plum curculio, Tarnished plant bug, Green fruit worm, Oriental fruit moth.	Apply when insects first appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears	Pear psylla, Green fruitworm and Codling moth.	Apply when needed. Do not apply more than 3 applications per season. Can be applied up to 14 days office harvest.

NUISANCE PESTS IN OUTDOOR AREAS

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs. When spraying plants, do not operate closer than, 18 % spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment exceed 90°F. inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not

OUTSIDE SURFACES OF I LDING

Use for control of Ants, Asian ladybugs, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

OTHER OUTDOOR AREAS

Use for control of Ants, Crickets, Earwigs, Fleas, Lone star ticks for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly.

For ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

Fire Ant Control Individual Mound and Plant Container Treatment

Gently apply over the surface of each mound. Thoroughly wet mound and surrounding area of 3 x 4 ft. (12 sq. ft.) For best results apply in cool weather 65°F-80°F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration, which will reduce product effectiveness. <u>Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.</u>

For use only as an aid in reducing annoyance from these insects on the following: Cluster flies, Flying moths, Gnats, House flies, Mosquitoes. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Manufactured by Bonide Products, Inc. 6301 Sutliff Road Oriskany, NY 13424

[Batch Code to be inserted at production]

©Bonide Products, Inc. All Rights Reserved

Optional Marketing Statements (Continued from page 1)

Protects your Houseplants, Roses, Flowers, Vegetables, Fruit, and Nuts from: Alfalfa Loopers, Ants, Aphids, Armyworms, Asian Ladybugs, Bagworms, Beetles, Borers, Budworms, Cabbageworms, Chinch Bugs, Cicadas, Citrus Black Flies, Codling Moths, Corn Earworms, Cutworms, Elm Spanworms, Fall Cankerworms, Green Fruitworms, Grasshoppers, Inchworms, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Lygus bags, Mealybugs, Mole Crickets, Naval Orange Worms, Scales, Sod Webworms, Sowbugs (pillbugs), Spider Mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies, and many others, of the control of t

Protects your home from: Carpenter Ants, Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Milipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clovermites, Cheese Mites, Granafy Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites, Cadelles, Fire Ants, Earwigs, Fleas, and Lone Star Ticks.

Use in and around the home on...Houseplants, Vegetables, Roses, Flowers, Ornamentals