

4-407

3-18-2004

1/10

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 4-407	2. EPA Product Manager La Rocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Bonide Eight Yard + Garden Ready to Spray	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Bonide Products, Inc. 6301 Sutliff Road Oriskany, NY 13424 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  MAR 18 2004
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
Notification of Additional Pest: Locust.  
see attached for placement

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, but no other changes have been made to the labeling or the CSF of this product. Bonide Products, Inc. understands that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA and further understands that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and Bonide Products, Inc. may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	If "Yes" Unit Packaging wgt. No. per container		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kate Entwistle	Title Registration specialist	Telephone No. (Include Area Code)
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped) 
2. Signature Kate Entwistle	3. Title Registration Specialist	
4. Typed Name kate Entwistle	5. Date 3-1-04	

2/10



# YARD & GARDEN READY TO SPRAY

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**OCT 21 2003**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
4-407

[Alternate Brand Names: Bonide Bug Beater II Yard & Garden Insect Control Ready to Spray; Tickmaster; Mosquito Beater Ready to Spray; Bonide Eight Insect Control Yard & Garden; Eight Yard & Garden]

Optional Marketing Statements:

Spray Around Shrubbery and Other Places Insects (Mosquitoes) Congregate

· NO MIX · NO MESS · NO FUSS

Easy to Apply

Easy to Use

Water Based

Will Not Stain

Contact and Residual Spray

For Residential Use Only

Kills Troublesome Outdoor Insects Which Destroy Lawns, Roses, Flowers, Trees and Shrubs, Vegetables, Fruit, and Nuts

Contains Permethrin, a Long Lasting Insecticide Which Provides Contact and Residual Control for up to 4 Weeks of Many Hard-to-Kill Sucking and Chewing Insects

Kills Over 100 Different Insects

Protects Your Lawns, Trees & Shrubs, Roses & Flowers, Fruits & Nuts, and Vegetables from: Alfalfa loopers,

(locust),

- Ants, Aphids, Armyworms, Asian Ladybug, Bagworms, Beetles, Borers, Brown dog ticks, Budworms,
- Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets,
- Cutworms, Earwigs, Elm spanworms, Fall cankerworms, Fire ants, Fleas, Green fruitworms, Gnats, Grasshoppers,
- Houseflies, Inchworms, Japanese beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs,
- Mealybugs, Millipedes, Earworms, Mole crickets, Mosquitoes, Moths, Navel orangeworms, Scales, Sod webworms, Sowbugs (Pillbugs), Spider mites, Spring cankerworms, Tent caterpillars, Thrips, Tobacco budworms,
- Weevils, Whiteflies and many others.

locust

**ACTIVE INGREDIENT:**

Permethrin (CAS #52645-53-1)..... 2.5%

**OTHER INGREDIENTS:**..... 97.5%

**TOTAL**..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See back/side/other panel(s) for additional Precautionary Statements.

**NET CONTENTS:** \_\_\_\_\_

3/10

<b>FIRST AID</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network at 1-800-858-7378 for emergency medical treatment information.	
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. Avoid contamination of ornamental fish ponds. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

**For Residential use only.**

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

**SHAKE WELL BEFORE USING**

41/10

**HOSE END SPRAYER: Easy to Use**

1. Make sure knob control lever is in the OFF position. Shake well and attach to hose.
2. Turn water on at faucet, aim nozzle toward lawn and turn knob control lever to ON to begin spraying.
3. When finished turn control lever to OFF position. Turn faucet to OFF position. Discharge residual water pressure by turning ON and OFF again.

Store out of reach of children.

People and pets should not be allowed in treated areas until spray has dried.

**DIRECTIONS FOR APPLYING**

**HOME LAWNS**

	<b>Pests:</b>	<b>How to Use:</b>
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine	Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Fleas, Grasshoppers, Japanese beetle grubs, Mole crickets, Sod webworms, Ticks that may carry Lyme Disease, Leafhoppers	Home lawns should be 3 inches or less in height at the time of application. 1 Quart of this concentrate as applied through the Hose-End sprayer treats 5,000 sq. ft. For heavy infestations, repeat application after 2 weeks.

**TREES & SHRUBS**

(Locust)

	<b>Pests:</b>	<b>How to Use:</b>
Arizona Cypress, Azalea, Birch, Cherry, Non-bearing Citrus, Conifers, Elm, English Ivy, Euonymous, Weeping fig, Fir, Honeysuckle, Lilac, Mock-orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar, Rhododendron, Taxus	Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Leafrollers, Locust, Mealybugs, Spider mites, Whiteflies, Gypsy Moth, Tent Caterpillar, Japanese Beetles	Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use at intervals of 4-8 days. Application can be made up to the day of collection.
Cherry, Honeysuckle	Fall webworms	
Conifer	Budworms	
Elm	Elm leaf beetles, Elm spanworms, Fall webworms	
Douglas and Grand Firs	Tussock moths	
Non-bearing Citrus	Citrus black flies	
Oak	Oakworms	
Pine	Pine beetles, Pine moths, Pine needleminers, Needle scales	

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

5/10

(Locust)

**ROSES & FLOWERS**

	<b>Pests:</b>	<b>How to Use:</b>
Ageratum, Aster, Ardisia, Azalea, Baby's breath, Begonia, Bird's-nest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Bagworms, Cabbage worms, Cicadas, Cornworms, Cutworms, Exposed thrips, Fall cankerworms, Fall webworms, Fungus gnats, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scale crawlers, Spring cankerworms, Spider mites, Whiteflies, Japanese Beetles, Rose Chafer, Psyllids, Scale Insects	Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50°F to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection.

Locust

**FRUIT & NUT TREES**

**GENERAL DIRECTIONS**

Apply when insects appear. Wet plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

	<b>Pests:</b>	<b>How to Use:</b>
Almonds, Pecans	Navel orangeworms, Peach twig borers	Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Green fruitworms, Aphids, Leafrollers (Oblique-banded, Redbanded), Plum curculio, Rosy apple aphids, Spotted tentiform leafminers, Tarnished plant bugs, White apple leafhoppers, Japanese Beetles	Repeat as required to maintain control. Do not make more than 3 applications. Do not apply after petal fall.
Peaches	Borers (Lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs	Do not apply within 7 days of harvest. Do not make more than 8 applications per season.

6/10

**FRUIT & NUT TREES (Cont.)**

	<b>Pests:</b>	<b>How to Use:</b>
Pears (dormant through delayed dormant)	Pear psylla	Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.
Pears (summer)	Codling Moths, Green fruitworms, Pear psylla, Aphids	Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

**VEGETABLES**

	<b>Pests:</b>	<b>How to Use:</b>
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Apply as needed to ensure uniform coverage. Do not make more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
Asparagus (post-harvest)	Larval stages of asparagus beetle, Tarnished plant bug, Lygus bugs and Adult Japanese beetle	Treat ferns or bush growth after spear harvest when insects are present.
Beans	Mexican bean beetle, Leafhoppers, Flea beetles, Corn earworm, Lygus and Stinkbugs	Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.
Celery - See Leafy Vegetables		
Cole Crops. Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworms, Cabbage Aphids and Loopers, Diamondback Moths, Imported cabbage worms	Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage.
Corn - See Sweet Corn		

7/10

**VEGETABLES (Cont.)**

	<b>Pests:</b>	<b>How to Use:</b>
<b>Cucurbits:</b> Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds: melons including hybrids such as Cantaloupe, Casaba, Crenshaw, Honeydew, Honey Balls, Mango melon, Muskmelon, Persian melon, Pumpkin, Summer & Winter Squash, Watermelon	Aphids, Cabbage looper, Cucumber beetle (adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant bugs, Squash bugs	Apply as needed to ensure uniform coverage. Can be applied up to the day of harvest.
<b>Eggplant</b>	Colorado potato beetles, Vegetable leafminers	Apply as needed to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.
<b>Horseradish</b> (Foliar application)	Imported crucifer weevils	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.
<b>Leafy Vegetables:</b> Amaranth, Arrugula, Celery, Celtuce, Chervil, Corn salad, Chrysanthemum (edible- leaved and garland), cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Okra, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard	Alfalfa looper, Aphids, Beet armyworm, Cabbage looper, Corn earworm, Cutworms, European corn borer, Fall armyworm, Green cloverworm, Southern armyworm, Tobacco budworm, Vegetable leaf miner	Apply ever 5 to 10 days as needed. Do not make more than 7 applications per crop. Can be applied up to one day prior to harvest.
<b>Melons - See Cucurbits</b>		
<b>Potatoes</b>	Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, Cutworms, European corn borer, Potato psyllid, Tarnish plant bug	Apply as needed. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.
<b>Peppers</b>	Vegetable leafminers, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer	Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.

8/10

**VEGETABLES (Cont.)**

	<b>Pests:</b>	<b>How to Use:</b>
Strawberries, Raspberries, Blueberries	Strawberry weevils, Aphids, Spider mites, Lygus bugs, Flea beetles, Omnivorous leafhopper, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leafroller, Weevils	Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
Sweet Corn	Corn earworm, European corn borer, Fall armyworm, Southern armyworm	Apply every 5 days and when needed, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomatoes	Beet and southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle, Cabbage looper	Apply as needed for uniform coverage. Can be applied up to day of harvest.

**NUISANCE PESTS IN OUTDOOR AREAS**

**GENERAL DIRECTIONS**

Activate Hose-End Sprayer and spray mist listed areas for thorough coverage.

**OUTSIDE SURFACES OF BUILDING**


Use **Bonide Eight® Insect Control Ready-To-Spray** on Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (pillbugs) to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

**OTHER OUTDOOR AREAS**

Use **Bonide Eight® Insect Control Ready-to-Spray** on Ants, Asian Ladybug, Crickets, Earwigs, Fleas, Ticks (including ticks that may carry Lyme Disease) for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For Ants, thoroughly wet hills and runways. Repeat application as reinfestations occur.

Locust

To reduce annoyance from: **Cluster flies, Flying moths, Gnats, House flies, Mosquitoes,** Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate. Repeat as necessary.





9/10

**STORAGE AND DISPOSAL**

**Storage:** Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

**Disposal:** *If empty* – Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.


**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by:

Bonide Products, Inc.

Oriskany, NY 13424



**Bonide Products, Inc.**

6301 Sutliff Road  
Oriskany, NY 13424

Telephone: 315-736-8231  
Fax: 315-736-7582  
Email: mattv@bonide.com

10/110

March 1, 2004

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
US Environmental Protection Agency  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202-4501

**RE: Bonide Eight Yard & Garden Ready to Spray  
EPA Reg. 4-407  
Notification of:**

*Add pest 'Locust; see attached labeling*

To Whom It May Concern:

Enclosed, please find a completed form 8570-1, Notification for the addition of the pest 'Locust' and labeling for the above-mentioned registration. I trust you will find everything in order.

Should you have any questions or need additional information, please do not hesitate to contact me at (315) 736-8231 ext. 230.

Sincerely,



Kate Entwistle  
Registration Specialist

Enc.

