

4-423

2/26/2003

Page 184

Bonide Rotenone Garden Dust or Spray

BONIDE

**Rotenone
Garden Dust or Spray**

HOME VEGETABLES, ORNAMENTALS, AND SMALL BERRIES

CONTAINS 1.5% ROTENONE

ACTIVE INGREDIENTS:

Rotenone	1.5%
Other Cube Resins	2.2%

INERT INGREDIENTS: 96.3%

TOTAL 100.00%

EPA REG. NO. 4-423

ACCEPTED
FEB 26 2003

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 Registered under
 4-423


EPA EST. NO. 4-NY-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements.

Net Weight:



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, absorbed through the skin, or inhaled. Avoid inhalation of dust or spray. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash thoroughly before reuse.

FIRST AID	
Have label with you when obtaining treatment advice.	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by poison control center or doctor.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off Contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center, doctor, or 1-800-858-7378 immediately for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center, doctor, or 1-800-8585-7378 immediately for treatment advice.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Bonide Rotenone Garden Dust or Spray is suitable for application as a dust or spray.

USE RESTRICTIONS: FOR USE ON HOME VEGETABLES, ORNAMENTALS, AND SMALL BERRIES SITES AND PESTS listed in the table below.

Do not apply to food crops within 1 day of harvest.

APPLICATION DIRECTIONS:

AS A DUST:

Mixing Directions:

Use as is. No mixing required.

Application Directions:

See table below for timing of application. Apply directly from the shaker top carton or from a hand powder duster. Cover all parts of the plant with thin protective film, including top and underside of leaves and around the stem. One pound treats about 1500 sq. ft.

AS A SPRAY:

Mixing Directions:

Mix 1 cup of product with enough water to make a thin paste. Mix thoroughly so paste is smooth and free of lumps. Then add enough water to make one gallon of spray.

Application Directions:

See table below for timing of application. Spray thoroughly, to point of runoff. Apply to upper and lower leaf surfaces and especially around the vine stem. Agitate sprayer to keep nozzle from plugging.

REMOVAL OF RESIDUE: Remove residue by washing or brushing.

REPEAT APPLICATION: Repeat at 5 to 7 day intervals or as needed, (especially new growth) until pests are under control.

Sites, Pests, and Timing of Applications for Rotenone 1.5% Dust		
Sites	Pests	Timing of Applications
Vegetables		
Beans	Mexican Bean Beetle	Apply as soon as eggs begin to hatch. Pay special attention to leaf undersides.
Cabbage, Cauliflower	Imported Cabbage Worm , Diamond Back Moth	Apply as soon as worms appear. Use dust application.
Potato and Eggplant	Colorado Potato Beetle larvae, Potato Flea Beetles	Apply as soon as pests appear.
Cucumber, Melon	Cucumber Beetles, potato Flea Beetles	Apply as soon as plants appear above ground.
Tomato	Potato Flea Beetles, Colorado Potato Beetle larvae	Apply 2 or 3 days after plants are set in the field..
Ornamental Plants and Berry Plants		
Shrubs, Flowers, and Raspberries	Japanese Beetle and Aphids	Apply as soon as pests appear.

STORAGE AND DISPOSAL

Storage: Store only in original container, in a dry place inaccessible to children and pets.

Disposal: If empty: Do not reuse empty container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or **1-800-CLEANUP** for disposal instructions.

Never place unused product down any indoor or outdoor drain.

NOTICE: Buyer assumes all risks of use, storage and handling this product not in strict accordance with directions given herewith.

For information on pesticide products, (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Manufactured by:
Bonide Products, Inc.
6301 Sutliff Rd.
Oriskany, NY 13424