

TRUSTED SINCE 1926
BONIDE®

NO ESCAPE™

TRUSTED SINCE 1926
BONIDE®

Slug & Snail Killer




- Patented formulation to attract and control both juvenile and adult snails and slugs
- Apply one pound to 1000 sq. ft.
- Use around gardens, flowers, ornamentals, lawns, fruits and vegetables

NO ESCAPE™

Slug & Snail Killer

ACTIVE INGREDIENT
Metaldehyde..... 0.13%
OTHER INGREDIENTS..... 99.87%
TOTAL..... 100.00%
EPA Est. No. 4-NY-1 EPA Reg. No. 4-442



Net Wt. 2.5 lbs. (1.135kg)

PULL TAB:
Open Here to
Reveal Shaker
Spout

NO ESCAPE Slug & Snail Killer

No Escape Slug & Snail Killer

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply directly on fruits and vegetables and use only on soil. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

HOMEOWNER USE DIRECTIONS - Apply Bonide No Escape Slug & Snail Killer during the daytime in the shade or during overcast skies. Preferred time is during the evening hours, after sprinkling water, or after a rain. Scatter the Bonide No Escape Slug & Snail Killer onto the surface of the soil around gardens, flowers, ornamental plants and lawn areas. Apply one pound of Bonide No Escape Slug & Snail Killer to each 1,000 sq. ft. For heavier infestation apply over a smaller area. Apply only to soil surface around plants. Do not apply to foliage or other plant parts. See directions for use around vegetables and fruits. Snails and Slugs are likely to infest at the start of the spring growing season. Act early before damage occurs. Treatment every two weeks should give good control. During periods of heavy moisture, rainfall or waterings, it may be necessary to apply Bonide No Escape Slug & Snail Killer more often.

ADDITIONAL DIRECTIONS

Bonide No Escape Slug & Snail Killer is a patented formulation to attract and control both juveniles and adult snails and slugs. Bonide No Escape Slug & Snail Killer is easy to apply and economical. One pound of Bonide No Escape Slug & Snail Killer will bait 1,000 square feet. Snails and Slugs may be observed on-the-spot with an oozy slime excretion trail. Snails and slugs will eat anytime of the day, when overcast sky, rainy days, in damp cool places, under wood piles and trash, around and on ornamental gardens, on flowers along fence lines and around bushes. Scatter Bonide No Escape Slug & Snail Killer onto the soil during wet or dry occasion. Water may be added before applying but is not necessary.

GREENHOUSES, HOT HOUSE AND LATH STRUCTURES: Place around the pots, plants, and flats. Keep bait from coming in contact with plants. Scatter at anytime when dry, wet, or after watering.

FRUIT AND VEGETABLES

To control snails and slugs around the following fruits and vegetables, apply on to the soil surface around the plants. Do not contaminate edible parts.

FRUITS:

Apples, Apricots, Avocados, Blackberries, Blueberries, Boysenberries, Caneberries, Cherries, Citrus, Grapes, Loganberries, Peaches, Pears, Plums, Raspberries, Strawberries.

VEGETABLES:

Artichoke, Asparagus, Beans, Beets, Black-eyed peas, Broccoli, Brussel Sprouts, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Collards, Corn, Cowpeas, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lettuce, Melons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, and Watermelons.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. **DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

IF SWALLOWED: Call physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger.

Do not induce vomiting or give anything by mouth to an unconscious person.

SEE ELSEWHERE ON LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin or swallowed. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated areas.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

1.) Long-sleeved shirt and long pants 2.) Waterproof gloves 3.) Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater.

NON-AGRICULTURAL USE REQUIREMENT

The requirement in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children, pets, and other unprotected persons out of treated areas until soil has dried.

Bonide Products, Inc.
Oriskany, NY 13424

