

4-459

01-18-2012

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Bonide Products, Inc  
c/o Rebecca Horton  
Registrations by Design  
PO Box 1019  
Salem, VA 24153

JAN 18 2012

Subject: Labeling and CSF Amendment  
DRAGON CAPTAN WETTABLE  
EPA Registration No. 4-459  
Decision No. 456456  
Submission Date: 9/19/11

Dear Ms Horton:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended to identify a new supplier on the basic CSF dated 9/19/2011 and to revise the label is acceptable, provided you make the following changes:

1. In the "Flowers and Ornamentals" directions for use, delete the clause "except when dose is specified in terms of volume of mixture per square foot area," as there are no applications on the label which require this wording. The sentence will end with "...to wet the surface of the soil."

Submit one copy of your final printed labeling before you release the product for shipment. A copy of the labeling stamped "Accepted with Comments" is enclosed for your records. If you have any questions, please contact Dominic Schuler at (703) 347-0260 or via email at [schuler.dominic@epa.gov](mailto:schuler.dominic@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Tony Kish".

Tony Kish  
Product Manager (22)  
Fungicides Branch  
Registration Division (7504P)

2  
7

[FRONT PANEL]



**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
JAN 18 2012**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.**

4-459

## CAPTAN WETTABLE

(Additional Brand Names: Captan – 50% WP Fruit & Ornamental; Bonide Captan – 50% WP Fruit & Ornamental; Trusted Since 1926 Bonide Captan – 50% WP Fruit & Ornamental)

### Optional Marketing Statements:

For control of certain fungus diseases of listed fruits, nuts and ornamentals, ~~grasses~~ and as a soil treatment for the control of certain damping-off diseases.

FOR USE ON APPLES, PEACHES, GRAPES, STRAWBERRIES AND OTHERS LISTED.

CONTROLS LISTED DISEASES ON AZALEAS, CARNATIONS, CHRYSANTHEMUMS, ROSES AND OTHERS LISTED.

### ACTIVE INGREDIENT:

*Captan .....	48.93%
Related Derivatives .....	1.07%
<b>OTHER INGREDIENTS:</b> .....	<b>50.00%</b>
<b>TOTAL</b> .....	<b>100.00%</b>

\*N-Trichloromethylthio-4cyclohexene-1,2-dicarboximide

**Keep Out Of Reach Of Children**

## DANGER

(See back/side/other panels for additional Precautionary Statements & First Aid)

**Net Weight 8 OZ. (.23 kg)  
Packed by weight, not volume**

**[BACK/SIDE/OTHER PANEL(S)]**

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water of 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.</p>	
<p><b>NOTE TO PHYSICIAN:</b> Probable mucosal damage may contraindicate the use of gastric lavage.</p>	

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER:** Corrosive. Causes irreversible eye damage. May be fatal if inhaled. Do not get in eyes or on clothing. Do not breathe dust. Prolonged or frequently repeated skin contact may cause allergic skin reaction in some individuals.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):** Wear long pants and long-sleeved shirt, shoes plus socks, goggles or face shield, and chemical resistant gloves made of any waterproof material. Remove contaminated clothing and separately launder before reuse. Wash gloves thoroughly with soap and water before removing.

**USER SAFETY RECOMMENDATIONS: Users should:** 1) Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

5  
17

**ENVIRONMENTAL HAZARDS:** This chemical is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### FOR OUTDOOR RESIDENTIAL USE ONLY.

Do not allow this product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

**ENTRY RESTRICTIONS:** Do not allow people or pets to enter the treated area until sprays have dried.

**COMPATIBILITY:** This product is compatible with most commonly used insecticides and fungicides, expect highly alkaline materials such as hydrated lime. Captan should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixture

Where foliage applications are directed, mix designated amount of material in a little water to make a thin paste, add to specified amount of water, and keep mixture agitated during application. Use a compression type or hose-end sprayer.

FRUIT & NUT CROPS	
Size:	Amount of Spray Needed Per Tree Diluted Spray:
Young or Dwarf	1 to 2 pints
Medium	1 to 2 quarts
Large	½ to 3 gallons

**ALMONDS:** *Anthracnose, brown rot, twig and blossom blight, shothole, scab, leaf blight* – Use 1 to 2½ Tbs. per gal. of water. Apply at popcorn, bloom and petal fall stages and full cover sprays. Do not harvest almonds within 30 days of last application.

**APPLES (Eastern U.S.):** *Primary scab, black rot (frog-eye), botrytis blossom-end-rot*– Use 1½ Tbs. per gal. of water. Apply in prebloom, bloom, petal fall and first cover sprays at 10 to 14 day intervals. *Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryosphaeria rot, bitter rot* – Use 1½ Tbs. per gal. of water. Apply in second and late cover sprays at 10 to 14 day intervals. May be applied up to harvest. **Note:** Do not apply this product with or closely following, or in alternation with wettable sulfur products on sulfur-sensitive varieties of apples such as Baldwin, King, Red Delicious, Staymen, etc. as severe injury and defoliation may occur.

**APPLES (Western U.S.):** *Primary scab* – Use 1 to 1½ Tbs. per gal. of water. For **secondary scab** in mid-summer coverage spray – Use 1 Tbs. per gal. of water. **(Pacific Northwest):** *Bull's eye rot, botrytis*

5  
7

rot -- Use 1 Tbs. per gal. of water. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest. May be applied up to harvest.

**APRICOTS:** *Brown rot (twig blight), jacket rot* – Use 1 to 1½ Tbs. per gal. of water. Apply in red bud, bloom, 75% petal fall and cover sprays. May be applied up to harvest.

**BLACKBERRIES, RASPBERRIES:** *Anthracnose, botrytis, spur blight* – Use 3 Tbs. per gal. of water. Apply when blossoms are in bud and young canes are 8 to 10 inches long. Make a second application 2 weeks later. Apply as a fall spray after old canes are removed. *Fruit rot* – Use 5 Tbs. per gal of water. Apply at early (5 to 10%) bloom and again at full bloom. Reapply at 10 to 14 days, as needed. Do not apply within 3 days of harvest.

**BLUEBERRIES (Eastern U.S.):** *Botrytis gray mold or berry rot, mummy berry* – Use 2 Tbs. per gal. of water. Start spray program when buds swell or when buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom. May be applied up to harvest.

**BLUEBERRIES (Western U.S.):** *Botrytis gray mold or berry rot, mummy berry* – Use 1 to 2 Tbs. per gal. of water. Begin at mid-bloom and repeat at 7-10 day intervals until maturity. May be applied up to harvest.

**CHERRIES (Eastern U.S.):** *Brown rot, leaf spot, botrytis rot, blossom blight* – Use 1½ Tbs. per gal. of water. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be needed to control *blossom blight* during bloom. Repeat applications at 7 to 20 day intervals. May be applied up to harvest. Postharvest Sprays: *Leaf spot* – Use 1½ Tbs per gal. of water. Apply immediately after harvest and again in 10 to 14 days.

**CHERRIES (Western U.S.):** *Brown rot blossom blight, brown rot (fruit), leaf spot* – Use 1 to 1½ Tbs. per gal. of water. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays. May be applied up to harvest.

**GRAPES (U.S. except CA):** *Phomopsis can and leaf spot, downy mildew, suppression of black rot* – Use 1 to 1½ Tbs. per gal. of water. Apply sufficient spray volume to provide thorough, even coverage when shoots are ½ to 1½ inches long, when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom, and continue at 10 to 14 day intervals as needed. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. May be applied up to harvest.

**GRAPES (in CA):** *Bunch rot (botrytis)* – Use 1½ Tbs. per gal. of water. Make 2 applications before bloom and immediately after bloom. Repeat periodically, making 3 cover applications before the bunches close. *Phomopsis cane and leaf spot (current season infection)* – Use 1 to 1½ Tbs. per gal. of water. Make first spray when green tissue begins to show, but before shoots are 6 to 8 inches long. May be applied up to harvest.

**NECTARINES:** *Brown rot, scab, blossom blight* – Use 1½ to 2½ Tbs per gal. of water. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control *blossom blight*. Repeat application at 7 to 14 day intervals as needed. *Coryneum blight (peach blight, shothole)* – Use 1½ to 2½ Tbs. per gal. of water. Apply in pink, bud, full bloom, petal fall, and cover sprays as necessary, and as a post harvest spray (before leaves drop). Preharvest spray may be applied up to day of harvest.

**PEACHES:** *Brown rot, scab, blossom blight* – Use 1 to 1½ Tbs per gal. of water. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control *blossom blight*. Repeat application at 7 to 14 day intervals as needed. *Coryneum blight (peach blight, shothole)* – Use 1½ Tbs. per gal. of water. Apply in pink, bud, full bloom, petal fall, and cover sprays as necessary, and as a post harvest spray (before leaves drop). Preharvest spray may be applied up to day of harvest.

**PLUMS, FRESH PRUNES (Eastern U.S.):** *Brown rot* – Use 1½ Tbs. per gal. of water. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed. Continue applications through harvest if conditions favor *brown rot*. May be applied up to harvest.

**PLUMS, FRESH PRUNES (Western U.S.):** *Brown rot* – Use 1 to 1½ Tbs. per gal. of water. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as needed. *Prune russet scab (lacy scab)* – Use 1 to 1½ Tbs. per gal. of water. Apply at full bloom. May be applied up to harvest.

**STRAWBERRIES:** *Botrytis rot (gray mold), leaf spot, fruit rot* – Use 4 to 8 Tbs. per gal. of water. Apply sufficient spray volume to provide thorough, even coverage. Begin applications when new growth starts in Spring and before fruit starts to form. Repeat at 7 to 14 day intervals. Under conditions favorable to fruit rot, continue applications through harvest, treating immediately after each picking. May be applied up to harvest.

**FLOWERS AND ORNAMENTALS**

**Use Precautions:** Do not apply to lawns. Do not apply to ornamental plants beyond the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot area.

Crop	Disease(s)	Rate per Gallon of Water
Azaleas	Damping-off of cuttings	3 Tbs.
	SPECIFIC DIRECTIONS: Dip cuttings before bedding.	
	Petal blight	1½ Tbs.
	SPECIFIC DIRECTIONS: Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.	

7  
-  
7

Begonias (Tuberous)	Damping-off, Tuber rot	4¾ Tbs.
	SPECIFIC DIRECTIONS: Dip tubers in diluted mixture for 30 minutes, drain and plant.	
Camellias	Petal blight	1 Tbs.
	SPECIFIC DIRECTIONS: Apply to drench soil around plants, beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.	
Carnations	Alternaria leaf spot, rust	1½ Tbs.
	SPECIFIC DIRECTIONS: Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews.	
	Damping-off of cuttings	3 Tbs.
	SPECIFIC DIRECTIONS: Dip cuttings before bedding.	
Chrysanthemums	Botrytis flower blight, septoria leaf spot	2 Tbs.
	SPECIFIC DIRECTIONS: Apply at first sign of disease. Repeat at 7 to 14 day intervals.	
	Damping-off of cuttings	3 Tbs.
	SPECIFIC DIRECTIONS: Dip cuttings before bedding.	
Gladiolus (Corms)	Corm rot and decay, damping-off	½ - 1¼ Tbs.
	SPECIFIC DIRECTIONS: Dip corms 25 minutes. Drain and plant.	
Roses	Black spot, botrytis blossom blight	2 Tbs.
	SPECIFIC DIRECTIONS: Apply at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.	

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place away from feed and foodstuffs.

**PESTICIDE DISPOSAL & CONTAINER HANDLING:** **If empty:** Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyers Guarantee Limited to Label Claims.

©Bonide Products, Inc. All Rights Reserved

Bonide Products, Inc.  
6301 Sutliff Road  
Oriskany, NY 13424