




Laboratories

700 MARIETTA STREET, N. W., ATLANTA, GEORGIA 30318



FRONT PANEL

SANICIDE

SANITIZER AND DISINFECTANT

ACTIVE INGREDIENTS:

Alkyl (C<sub>12</sub> 61%, C<sub>14</sub> 23%, C<sub>16</sub> 11%, C<sub>8</sub>, C<sub>10</sub>  
and C<sub>18</sub> 5%) Dimethyl Benzyl Ammonium Chloride ..... 10.25%

INERT INGREDIENTS:

Water ..... 89.75%


Phenol Coefficients

S. typhosa ..... 28  
S. aureus ..... 40

This product fulfills the criteria of Appendix F as revised March 12, 1956 of the Milk ordinance and code of the U.S. Public Health Service in waters up to 550 ppm of hardness, calculated as CaCO<sub>3</sub> when tested by the method outlined by Chambers.

USDA Registration NO. 5-d

Keep out of reach of children. Sanicide is a concentrated disinfectant. Avoid contact with skin and eyes, in case of contact flush with plenty of water and get medical attention. Harmful if swallowed. Avoid contamination of food.



Net Contents

EMPIRE LABORATORIES

700 MARIETTA S. N. W.  
ATLANTA, GA.

DIRECTIONS FOR USE IN EATING AND DRINKING ESTABLISHMENTS

1. Scrape and prewash dishes, glasses, utensils and silverware.
2. Wash with suitable detergent.
3. Rinse with clean water.
4. Sanitize by immersing for two minutes in a 200 ppm solution obtained by adding one ounce of Sanicide in each 3.75 gallons of water.
5. Place sanitized glasses, dishes, silver and utensils on a drain board or rack to air dry.

IN DAIRY FARMS, DAIRIES AND FOOD PROCESSING PLANTS

1. Clean equipment and utensils using a suitable detergent.
2. Sanitize by immersing, spraying, mopping or circulating a 200 ppm solution of Sanicide obtained by adding one ounce of Sanicide to each 3.75 gallons of water.
3. Rinse with clear water before allowing sanitized surface to contact with food.

BACK PANEL

Directions for use as a disinfectant in hospitals, homes, public buildings, schools and institutions.

First clean surface with suitable detergent, then wipe down, spray or mop with a 400 ppm solution of Sanicide obtained by adding two ounces of Sanicide to each 375 gallons of water.

In order to expedite registration of your product send five copies of the proposed label to U.S.D.A., Pesticide Registration Section, Washington, D.C. 20250 referring to U.S.D.A. registration no. 2823-2 stating the active ingredient in your product is Roccal 50% manufactured by Sterwin Chemicals, New York, New York.