

D I R E C T I O N S

Remove leaves, dead or tall grass and other debris and mark off the area to be treated (ill. 1,000 sq. ft. i.e. 25'x40'). Follow the directions for the insect to be controlled applying the correct rate or amount per 1,000 sq. ft.

LAWN CHINCH BUGS: Thoroughly water grass just before application (25 gals./1,000 sq.ft.). Apply at rate of 1 lb. per 1,000 sq. ft. (25 lbs./25,000 sq. ft.). Do not water lawn for 1-2 days after application. For longer residual control and in lawns of thicker growth (i.e. St. Augustine), apply at rate of 2 lbs. per 1,000 sq. ft. (25 lbs./12,500 sq. ft.). A second application in 7-10 days may be necessary.

SOD WEBWORMS (Lawn Moths), ANTS, SOWBUGS, COLLEMBOLA (Springtails), ARMYWORMS, CUTWORMS, EARWIGS, CRICKETS, CLOVER MITES, LAWN BILLBUGS: Water lawn lightly (3 gals./1,000 sq.ft.), or apply when dew is present. Apply at rate of 1 lb. per 1,000 sq. ft. (25 lbs./20,000 sq. ft.).

For Lawn Billbugs: Irrigate the lawn thoroughly after treatment. For ants, brown dog ticks, clover mites, crickets, earwigs, fleas, millipedes, sowbugs and collembola (springtails), commonly found near the foundation of, and which may enter the home, treat a 5 ft. width up to and around the foundation.

FLEAS, TICKS, BROWN DOG TICKS, CHIGGERS, LEANHOOPPERS, BERMUDA MITES: Water lawn lightly (3 gals./1,000 sq.ft.). Apply at rate of 1 lb. per 1,000 sq.ft. (50 lbs./50,000 sq.ft.). For Brown Dog Ticks do not apply to animals; treat areas under shrubbery and grass near house in particular.

MILLIPEDS: Water lawn lightly (3 gals./1,000 sq.ft.). Apply at rate of 2 1/2 lbs. per 1,000 sq. ft. (25 lbs./10,000 sq.ft.).

GREEN JUNE BEETLE, WHITE GRUBS (Japanese Beetle, Western and European Chafer Grubs): Apply at rate of 2-3 1/4 lbs. per 1,000 sq. ft. (25 lbs./3,900 sq.ft.). Water heavily (100 gals./1,000 sq.ft.) to move active ingredients into root zone after application.

ANTWORMS: Apply at rate of 3/4 lbs. per 1,000 sq.ft. (25 lbs./33,000 sq.ft.). Water into top 2 to 4 inches of soil (10-15 gals./1,000 sq.ft.).

CRICKETS: Apply at rate of 1 lb. per 1,000 sq. ft. (25 lbs./25,000 sq. ft.). Water into top 1-2 inches of soil (10-15 gals./1,000 sq. ft.).

SUGGESTED SPREADER SETTINGS: These settings will give approximate starting points for spreader application. In a small measured area with a weighed quantity and adjust your spreader by opening or closing to obtain correct coverage. Adjustments will be necessary due to condition of spreader, rate of walk and terrain characteristics.

SPREADER	ACCEPTED RATE		
	1 lb./1,000 sq. ft.	2 1/2 lbs./10,000 sq. ft.	6 lbs./25,000 sq. ft.
Lawn Beauty/Agrow	1/2	1	2
Modern/Scotts	1/2	1	2
Jackson/Patch	1/2	1	2
Byline/Ortho	1/2	1	2