

PM 04

16-12

8/1/62

Pg 175

Best Available Image

FRONT PANEL

New! 4 lbs. New Wt.

BLUE DRAGON

Contains -

SEVIN*

* Trade Mark of UCC for
1-naphthyl N-methylcarbamate

Kills Japanese beetles, grasshoppers, Mexican
bean beetle, locusts, ~~leafhopper~~ ^(cicada), tomato Hornworm,
corn earworm, flea beetle, cabbage looper and
many other insects attacking vegetables, fruits,
ornamentals, shrubs, ~~and lawns~~.

ANALYSIS

ACTIVE INGREDIENTS

1-naphthyl N-methylcarbamate 2%

Sulphur 10%

INERT INGREDIENTS 88%

J. M. HARRIS & CO., INC. - ROANOKE, VIRGINIA

ACCEPTED
AUG 1 1962
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 16-12

SIDE PANEL

New!

BLUE DRAGON

New Blue Dragon is safe to any domestic animal, pet or human when used as directed.

CAUTION:

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed or food stuffs. Keep out of reach of children.

NOTICE: Sold without warranty as to hazards or results.

BACK PANEL

New!

BLUE DRAGON

GENERAL INFORMATION

New BLUE DRAGON is an outstanding, new, effective insecticide that controls a wide range of insects and has ability to control insects resistant to other insecticides. Apply when insects appear, before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces; if needed, repeat every 7-10 days. Do not use with lime or other alkaline material. New Blue Dragon is one of the safest insecticides available and may be applied up to harvest except when noted.

DIRECTIONS

BEANS - For control of Mexican bean beetle, leafhopper, fleabeetles, Japanese beetle, velvet bean caterpillar, apply 3/4 lbs. per 1,000 sq. ft. For control of lygus, stink bug, corn earworm, tarnished plant bug, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.

CORN - For control of Corn earworm, European corn borer, Japanese beetle, apply 1 1/4 to 2 1/2 lbs. per 1,000 sq. ft. Allow 7-days before harvest or grazing of forage.

4/5

BACK PANEL - Cont'd.

POTATOES, TOMATOES, EGGPLANT and PEPPERS - For control of Colorado potato beetle, flea beetles, leafhoppers, apply $3/4$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For control of tomato fruitworm, fall army worm, tomato hornworm, European corn borer, stink bugs, lace bugs, lygus bugs, apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, watermelon, honey dew) - For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles, leafhoppers, apply $3/4$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft.

LETTUCE, CARROTS - For control of flea beetles, leafhoppers, harlequin bug, apply $3/4$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. - for sixspotted leafhopper apply $1\frac{1}{4}$ to $1-3/4$ lbs. per 1,000 sq. ft. - for corn earworms, army worms, cabbage caterpillars, spittlebugs, lygus and stink bugs, apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft. Do not apply to lettuce within 3 days of harvest.

ASPARAGUS - For control of asparagus beetle on spears, apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft.; for asparagus beetle, apache cicada on ferns and brush, apply $2\frac{1}{2}$ to 4 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

OKRA, CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, KOHLRABI - For control of flea beetle, harlequin bug, apply $3/4$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft.; for corn earworm, stink bug army worms, cabbage caterpillars, apply $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

75

ACCEPTED
AUG 1 1962
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 16-17

BACK PANEL - Cont'd.

ORNAMENTALS, FLOWERS, SHRUBS and SHADE TREES - For control of Japanese beetles, ^{grasshoppers,} locust, ^(Cicada) bagworm, lace bugs, leafhoppers, scale insects, rose aphid, rose slug, tent caterpillar, gypsy moth, boxwood ^{leaf-miner} and many others, apply 2 to 2 1/2 lbs. per 1,000 sq. ft.

GRAPES - For control of leafhopper, Japanese beetle, grape berry moth, June beetles, apply 3/4 to ~~1 3/4~~ lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of meadow spittle bug, leaf roller, strawberry weevil, apply 1 1/2 to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

TOBACCO PLANT BEDS - For control of tobacco flea beetle, green June beetle grubs. *Do not apply to plants in flower. Plant, apply as needed at 12 to 14 days intervals and repeat transplanting.*

POULTRY - PIGEONS - For control of northern fowl mite, *pub lice* and chicken mite, fleas, ~~lice~~ on birds, apply 2 1/2 lbs. per 100 birds; for chicken mite, fleas, bedbugs, apply 2 1/2 lbs. per 40 sq. ft. of floor litter, *and to roosts and adjacent areas. Force dust into cracks and crevices. Repeat as necessary.*

ANIMALS, DOGS, CATS - For control of fleas, brown dog tick, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens under 4 weeks old.

Do not apply directly to eggs or nest litter. Do not contaminate food or drinking water. Do not leave birds in treated houses within 7 days of application.