

PM 04

11/4/63

16-12

pg 1/4

FRONT PA

New

1 LB. NET WT.

BLUE DRAGON

Contains

Sevin*

*Trade Mark of UCC for
1-naphthyl N-methylcarbamate

BLUE DRAGON

ACCEPTED

NOV 4 1963

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 16-12

Kills Japanese beetles, Mexican bean beetle, tomato hornworm,
corn earworm, flea beetle and many other insects attacking veg-
etables, fruits, ornamentals and shrubs.

ANALYSIS

ACTIVE INGREDIENTS

1-naphthyl n-methylcarbamate 2%

Sulphur 10%

INERT INGREDIENTS 88%

Manufactured By

J. M. HARRIS & CO., INC.

ROANOKE, VA.

DO NOT REVERSE APPLICATION

2/4

SIDE PANEL

New

BLUE DRAGON

New Blue Dragon is one of the safest insecticides available and may be applied up to harvest except when noted.

CAUTION

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and food stuffs. Keep out of reach of children.

NOTICE: Sold without warranty as to hazards or results.

DATE APPLICATION OCT 29 1953



NEW
BLUE DRAGON

General Information

New BLUE DRAGON is an outstanding, new, effective insecticide that controls a wide range of insects and has ability to control ^{INSECTS} resistant to other insecticides. Apply when insects appear, before damage from feeding occurs. Dust thoroughly to upper and lower leaf surface; if needed, repeat every 7-10 days. Do not use with lime or other alkaline material. New Blue Dragon is one of the safest insecticides available and may be applied up to harvest except when noted.

DIRECTIONS

BEANS-For control of Mexican bean beetle, leafhopper, flea beetles, Japanese beetle, velvet bean caterpillar, apply 3/4 lb. per 1,000 sq. ft. For control of lygus, stink bug, corn earworm, tarnished plant bug, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.


CORN-For control of Corn earworm, European corn borer, Japanese beetle, apply 1 1/4 to 2 lbs. per 1,000 sq. ft. Allow 7 days before harvest or grazing of forage.

POTATOES, TOMATOES, EGGPLANT and PEPPERS-For control of Colorado potato beetle, flea beetles, leafhoppers, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft. For control of tomato fruitworm, fall army worm, tomato hornworm, European corn borer, stink bugs, lace bugs, lygus bugs, apply 1 1/4 to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Canteloupe, watermelon, honey dew) For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles, leafhoppers, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH-For control of flea beetles, leafhoppers, harlequin bug, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft. For spotted leafhoppers, corn earworm, army worm, cabbage caterpillar, spittlebugs, lygus and stink bugs apply 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (Except head lettuce 3 days)

ASPARAGUS-For control of asparagus beetle on spears, apply 1 1/4 to 2 1/2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest. *APPLY during cutting season. within 3 days. Do not repeat application.*



CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS-For control of flea beetles, harlequin bugs, apply 3/4 to 1 1/4 lbs. per 1,000 sq. ft. For corn earworm, stink bug, army worm, cabbage caterpillar apply 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEA and other ORNAMENTALS-For control of Japanese beetle, bag worm, flea beetles, leafhoppers, scale insects, tent caterpillar and many others, apply 2 to 2 1/4 lbs. per 1,000 sq. ft.

GRAPES-For control of leafhoppers, Japanese beetle, grape berry moth, June beetles, ~~Black Rot, Downy and Powdery Mildew~~, apply 3/4 to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES-For control of meadow spittle bug, leaf roller, strawberry weevil, apply 1 1/4 to 2 lbs. per 1,000 sq. ft. *Do not apply within 1-day of harvest*

ANIMALS, DOGS, CATS-For control of fleas, brown dog tick, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens under 4 weeks old.

APPLICATION 222 8 5 700

