

PM 04

16-12

12/19/66

P 174

November 4, 1963 (original)
September 28, 1966 (revised)

USDA Reg. No. 16-12

BLUE DRAGON

ACCEPTED
DEC 19 1966
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 16-12 SUBJECT
TO ATTACHED COMMENTS.



24

FRONT PANEL

BLUE DRAGON
Contains 2% SEVIN*

Net Weight
1 lb., 2 lb., 4 lb.
& 25 lbs.

**KILLS JAPANESE BEETLE, MEXICAN BEAN BEETLE, TOMATO HORNWORM,
CORN EARWORM, FLEA BEETLE AND MANY OTHER INSECTS ATTACKING
VEGETABLES, FRUITS, ORNAMENTALS AND SHRUBS.**

ACTIVE INGREDIENTS	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate).....	2%
Sulfur.....	10%
INERT INGREDIENTS.....	88%
TOTAL	100%

CAUTION: Keep out of reach of children. See back/side panels
for additional cautions.

*Trade Mark of UCC for 1-naphthyl N-methylcarbamate

DRAGON CHEMICAL CORPORATION
ROANOKE, VIRGINIA
USDA Reg. No. 16-12

Inert Ingredient: Talc



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON is an outstanding, new, effective insecticide that controls a wide range of insects and has ability to control insects resistant to other insecticides. Apply when insects appear, before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces; if needed, repeat every 7-10 days. Do not use with lime or other alkaline material. Blue Dragon may be applied up to harvest except where noted.

DIRECTIONS

BEANS - For control of MEXICAN BEAN BEETLE, LEAFHOPPER, FLEA BEETLES, JAPANESE BEETLE, VELVET BEAN CATERPILLAR, apply $\frac{3}{4}$ lb. per 1,000 sq. ft. For control of LYGUS, STINK BUG, CORN EARWORM, TARNISHED PLANT BUG, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft.

CORN - For control of CORN EARWORM, EUROPEAN CORN BORER, JAPANESE BEETLE, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft. Allow 7 days before harvest or grazing of forage.

POTATOES, TOMATOES, EGGPLANT and PEPPERS - For control of COLORADO POTATO BEETLE, FLEA BEETLES, LEAFHOPPERS, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For control of TOMATO FRUITWORM, FALL ARMY WORM, TOMATO HORNWORM, EUROPEAN CORN BORER, STINK BUGS, LACE BUGS, LYGUS BUGS, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (CANTELOUPE, WATERMELON, HONEY DEW) - For control of PICKLEWORM, MELONWORM, CUCUMBER BEETLES, SQUASH BUGS, FLEA BEETLES, LEAFHOPPERS, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIP GREENS, MUSTARD GREENS, SPINACH - For control of FLEA BEETLES, LEAFHOPPERS, HARLEQUIN BUG, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For SIXSPOTTED LEAFHOPPERS, CORN EARWORM, ARMY WORM, CABBAGE CATERPILLAR, SPITTLEBUGS, LYGUS and STINK BUGS, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce 3 days).

ASPARAGUS - For control of ASPARAGUS BEETLE on spears, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS - For control of FLEA BEETLES, HARLEQUIN BUGS, apply $\frac{3}{4}$ to $1\frac{1}{4}$ lbs. per 1,000 sq. ft. For CORN EARWORM, STINK BUG, ARMY WORM, CABBAGE CATERPILLAR, apply $1\frac{1}{2}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.



4/4

ROSES, AZALEA and Other ORNAMENTALS - For control of JAPANESE BEETLE, BAG WORM, FLEA BEETLES, LEAFHOPPERS, SCALE INSECTS, TENT CATERPILLAR and many others, apply 2 to 2½ lbs. per 1,000 sq. ft.

GRAPES - For control of LEAFHOPPERS, JAPANESE BEETLE, GRAPE BERRY MOTH, JUNE BEETLES, apply ¾ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of MEADOW SPITTLEBUG, LEAF ROLLER, STRAWBERRY WEEVIL, apply 1½ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

ANIMALS, DOGS, CATS - For control of FLEAS, BROWN DOG TICK, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens under 4 weeks old.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

NOTICE

This material is sold without warranty as to hazards or results.

DRAGON CHEMICAL CORPORATION
ROANOKE, VIRGINIA

