

PM 04 16-12

5/29/69

17/4

acceptance steps on p. 2

FRONT PANEL

BLUE DRAGON GARDEN DUST

NET WEIGHT

CONTAINS: 2% Sevin and 10% Sulfur

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and Certain other insects.

ACTIVE INGREDIENTS

BY WEIGHT

\*Carbaryl (1-Naphthyl N-methylcarbamate)  
Sulfur

2%  
10%

INERT INGREDIENTS

88%

TOTAL:

100%

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional cautions.

USDA REG. NO. 16-12

\*Trade Mark of UCC for 1-naphthyl N-methylcarbamate

DRAGON CHEMICAL CORPORATION

ROANOKE, VIRGINIA

INERT INGREDIENT: TALC

2500

82-18

2/4

May 19, 1969

USDA REG. NO. 16-12

ACCEPTED  
MAY 29 1969  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 16-12

BLUE DRAGON GARDEN DUST

2500  
002-10

BACK PANEL

BLUE DRAGON GARDEN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS: AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

Blue Dragon Garden Dust is a ready to use combination of 2% Sevin and 10% Sulfur. Sevin controls certain insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear, before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces; if needed, repeat every 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted. Sulfur may burn foliage when temperatures are excessively high.

DIRECTIONS

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply 3/4 lb. per 1,000 sq. ft. For control of Lygus, Stink Bug, Corn Earworm, Tarnish Plant Bug, apply 1-1/4 to 2 lbs. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN- For control of Corn Earworm, European Corn Borer, Japanese Beetle, apply 1-1/4 to 2 lbs. per 1,000 sq. ft. For Corn Earworm, apply to silks as soon as they appear. Make 3-4 applications at 2-3 day intervals. Re-apply immediately after rain. For Corn Borer, make one or more applications beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is achieved. Allow 7\* days before harvest or grazing of forage.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS- For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply 3/4 to 1-1/4 lbs. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, apply 1-1/4 to 2 lbs. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew)- For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply 3/4 to 1-1/4 lbs. per 1,000 sq. ft. Do not use sulfur on cucumbers, melons, or squash unless local experience has shown that sulfur will not burn the beans. *beans*

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH- For control of Flea Beetles, Leafhoppers, Harlequin Bug, apply 3/4 to 1-1/4 lbs. per 1,000 sq. ft. For six-spotted Leafhoppers, Corn Earworm, Armyworm, Cabbage Caterpillar, Spittlebugs, Lygus and Stink Bugs, apply 1-1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce - 3 days).

BACK PANEL

4/4

BLUE DRAGON GARDEN DUST

ASPARAGUS- For control of Asparagus Beetle on spears, apply 1-1/4 to 2-1/2 lbs. per 1,000 sq. ft. Do Not repeat application within 3 days. Do Not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSEL SPROUTS- For control of Flea Beetles, Harlequin Bugs apply 3/4 to 1-1/4 lbs. per 1,000 sq.ft. For Corn Earworm, Stink Bug, Armyworm, Cabbage Caterpillar, apply 1-1/4 to 2 lbs. per 1,000 sq. ft. Do Not apply within 3 days of harvest.

ROSES, AZALEA, AND OTHER ORNAMENTALS- For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillar and certain others, apply 2 to 1-1/4 lbs. per 1,000 sq. ft.

GRAPES- For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles apply 3/4 to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES- For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply 1-1/4 to 2 lbs. per 1,000 sq. ft. Do Not apply within 1 day of harvest.

DOGS, CATS- For control of Fleas, Brown Dog Tick, apply liberally and rub into skin, feet and legs, Dust sleeping quarters and infested areas weekly. DO NOT treat kittens under 4 weeks old.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE DUST. DO NOT TAKE INTERNALLY. Skin contact may harmful. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or feedstuffs. For protection of honey bees avoid use, if possible, during periods when honeybees are visiting crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

NOTICE

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS AND CAUTIONS STATED ON THIS LABEL.

DRAGON CHEMICAL CORPORATION

ROANOKE, VIRGINIA

