

16-12

07/05/1972


1/5

April 18, 1972

EPA Reg. No. 16-12

~~JUL 07 1972~~

BLUE DRAGON GARDEN DUST



FRONT PANEL

NET WEIGHT  
1, 2, 4, & 25 lb.

~~CONFIDENTIAL~~

**KILLS:** Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects.

<b>ACTIVE INGREDIENTS</b>		<b>By Wt.</b>
*Carbaryl (1-Naphthyl N-methylcarbamate).....		2%
<b>INERT INGREDIENTS</b>		98%
<del>Water</del> .....		<del>10%</del>
<del>etc.</del> .....		<del>2%</del>
	<b>Total</b>	<b>100%</b>

**CAUTION:** KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional cautions.

EPA Reg. No. 16-12

\*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

**ACCEPTED**

JUL 05 1972

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 16-12 SUBJECT  
TO ATTACHED COMMENTS.

Manufactured By  
**DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA**

BACK PANEL

BLUE DRAGON GARDEN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON GARDEN DUST ~~controls certain listed insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear, before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces, if needed, repeat every 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted.~~

DIRECTIONS

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1-1/4 lb. per 1,000 sq. ft. For control of Lygus, Stink Bugs, Corn Earworm, Tarnish Plant Bug, apply approximately 1-1/4 lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN - Use approximately 1-1/4 lb. to 2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, Flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS - For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1-1/4 lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, apply approximately 1-1/4 lb. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew) - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1-1/4 lb. per 1,000 sq. ft. ~~Some~~

~~and for cucumbers, pumpkins, squash unless local experience indicates otherwise.~~

ACCEPTED  
JUL 05 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 16-12 SUBJECT  
TO ATTACHED COMMENTS.

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH - For control of Flea Beetles, Leafhoppers, Harlequin Bug, apply approximately 3/4 to 1-1/4 lb. per 1,000 sq. ft. For six-spotted Leafhoppers, Corn Earworm, Armyworm, Cabbage Caterpillar and Stink Bugs, apply approximately 1-1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce - 3 days).

ASPARAGUS - For control of Asparagus Beetle on spears, apply approximately 1-1/4 to 2-1/2 lbs. per 1,000 sq. ft. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSEL SPROUTS - For control of Flea Beetles, Harlequin Bugs, apply approximately 3/4 to 1-1/4 lb. per 1,000 sq.ft. For Corn Earworm, Stink Bug, Armyworm, apply approximately 1-1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEAS, AND OTHER ORNAMENTALS - For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillars apply approximately 1-1/4 to 2 lbs. per 1,000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles, apply approximately 1-1/4 to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply approximately 1-1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

DOGS, CATS - For control of Fleas, Brown Dog Ticks, apply liberally and run into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens or puppies under 4 weeks old.

DRAGON CHEMICAL CORPORATION

Roanoke, Virginia 24019

ACCEPTED  
JUL 05 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 6-2 SUBJECT  
TO ATTACHED COMMENTS.

SIDE PANEL

BLUE  
DRAGON

CAUTION

HARMFUL IF INHALED OR SWALLOWED. DO NOT breathe dust. DO NOT take internally. Skin contact may be harmful. Wash hands before eating. Take shower or bath. after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

**ACCEPTED**  
JUL 05 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 16-12 SUBJECT  
TO ATTACHED COMMENTS.

