

ACCEPTED

May 8, 1973

UNL. THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 16-12-1 SUBJECT

16-12

4 LBS.



NET WT.



Blue Dragon

Blue Dragon

Blue Dragon



GARDEN DUST

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects.

ACTIVE INGREDIENT	BY WEIGHT
Carbaryl (1-Naphthyl N-methylcarbamate)	2%
INERT INGREDIENTS	98%
TOTAL	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional cautions.

EPA REG. NO. 16-12

Trade Mark of UCC for 1-naphthyl N-methylcarbamate

Manufactured By

DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24019

BLUE DRAGON GARDEN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON GARDEN DUST contains certain listed insects which attack vegetables, fruits, shrubs and trees. Apply when insects first appear, before damage from feeding occurs. Dust thoroughly to cover and force dust into crevices. If needed, repeat every 7 to 10 days. Do not mix with liquid or other dusting materials. Blue Dragon may be applied up to harvest except where noted.

DIRECTIONS

BEANS—For control of Mexican Bean Beetle, European Flea Beetle, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. Black Bean, Green Legume, French Bean, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. Do not feed treated beans to dairy or meat animals.

CORN—For control of European Corn Borer, Corn Earworm, and other insects, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For corn in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain. For silage, apply to the silage. For grain, apply to the grain.

POTATOES—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For potatoes in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

PEPPERS—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For peppers in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

SPINACH—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For spinach in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

LETTUCE—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For lettuce in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

AMARANTH—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For amaranth in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

STRAWBERRIES—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For strawberries in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

GRAPES—For control of Colorado Potato Beetle, apply approximately 1 lb. to 2 lbs. per 1,000 sq. ft. for outdoor use. For grapes in rows, apply to the base of the plant and the leaves. For silage, apply to the silage. For grain, apply to the grain.

DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24019

BLUE DRAGON



Best Copy

PM 04

16-12

5/8/73

P3181