



BLUE DRAGON®

BLUE DRAGON®

ACCIDENT
JUL 11 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 16-12



CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. **DO NOT** take internally. Wear regular long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

GARDEN DUST

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

KILLS: Japanese Beetles, Mexican Bean Beetles; Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects. **CONTAINS SEVIN**®*

ACTIVE INGREDIENT	BY WEIGHT
Carbaryl (1-Naphthyl N-methylcarbamate)	2%
INERT INGREDIENTS	98%
TOTAL	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN. See back/side panels for additional cautions.

SEVIN is a registered trademark of Union Carbide Corporation.

NET WEIGHT 4 LB.

PM04

16-12

7/11/82

PM 1/2



BLUE DRAGON GARDEN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON GARDEN DUST controls certain listed insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces, if needed, repeat 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

DIRECTIONS

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BEANS—For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1 1/4 lb. per 1,000 sq. ft. For control of Lygus, Slink Bugs, Corn Earworm, Tarnish Plant Bug, apply approximately 1 1/4 lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN—Use approximately 1 1/4 lb. to 2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the basal region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS—For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Flowerworm, European Corn Borer, Slink Bugs, Flea Beetles, apply approximately 1 1/4 lb. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew)—For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH—For control of Flea Beetles, Leafhoppers, Colorado Bug, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For six spotted Leafhoppers, Citrus Leafminer, Armyworm, and Slink Bugs, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce—3 days).

ASPARAGUS—For control of Asparagus Beetle on spears, apply approximately 1 1/4 to 2 1/2 lbs. per 1,000 sq. ft. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS—For control of Flea Beetles, Harlequin Bugs, apply approximately 3/4 to 1 1/4 lb. per 1,000 sq. ft. For Corn Earworm, Slink Bug, Armyworm, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEAS, AND OTHER ORNAMENTALS—For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillar, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft.

GRAPES—For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles; apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES—For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply approximately 1 1/4 to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

DOGS, CATS—For control of Fleas, Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens or puppies under 4 weeks old. Do not use on pregnant dogs.

STORAGE AND DISPOSAL: **Storage:** Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs. **Pesticide disposal:** Securely wrap original container in several layers of newspaper and discard in trash. **Container disposal:** Do not reuse empty bag. Discard bag in trash.



GARDEN DUST

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

EPA REG. NO. 16-12

EPA Est. 16-VA-1

Manufactured By
DRAGON CHEMICAL CORPORATION, ROANOKE, VIRGINIA 24019

RED
BLU

2/2