

PM 19 16-12

Please read instructions on reverse before completing form.

Form Approved OMB No 2070-0060. Approval expires 11-30-93

(A)  EPA	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number 188523
	Application for Pesticide:		

Section I

1. Company/Product Number 16-12	2. EPA Product Manager Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Blue Dragon^(R) Garden Dust	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Dragon Corporation P. O. Box 7311 Roanoke, VA 24019 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Compliance with PR Notice 93-3 -- Labeling Statement Prohibiting Application to Water

NOTIFICATION LABEL NOT REVIEWED PER PR NOTICE 93-3
DATE: 7/12/93

Section III

1. Material This Product Will Be Packaged In.				2. Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
	If "Yes," Unit Package wgt.	No. per container	If "Yes," Package wgt.	No. per container	<input checked="" type="checkbox"/> Paper
* Certification must be submitted.				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container 4 LB		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed To Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other (_____)	

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Rebecca M. Horton	Title Executive Assistant	Telephone No. (Include Area Code) (703) 362-2657
2. Signature <i>Rebecca M. Horton</i>		6. Date Application Received (Stamped)
3. Title Executive Assistant		
4. Typed Name Rebecca M. Horton		
5. Date 6/21/93		

NOTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 00-0
DATE 7/12/93

DRAGON®
BLUE DRAGON® GARDEN DUST

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects

Contains SEVIN® brand carbaryl insecticide

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 2%

INERT INGREDIENTS: 98%

TOTAL: 100%

Keep Out of Reach of Children
CAUTION
See back/side panels for additional cautions.

NET WEIGHT 4 LB

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON® GARDEN DUST controls certain listed insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces, if needed, repeat 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1 1/4 lb. per 1,000 sq. ft. For control of Lygus, Stink Bugs, Corn Earworm, Tarnish Plant Bug, apply approximately 1 1/4 lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN - Use approximately 1 1/4 lb. to 2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects

attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS - For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply approximately $\frac{3}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, apply approximately $1\frac{1}{4}$ lb. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew) - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply approximately $\frac{3}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH - For control of Flea Beetles, Leafhoppers, Harlequin Bug, apply approximately $\frac{3}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For six-spotted Leafhoppers, Corn Earworm, Armyworm and Stink Bugs, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce - 3 days).

ASPARAGUS - For control of Asparagus Beetle on spears, apply approximately $1\frac{1}{4}$ to $2\frac{1}{2}$ lbs. per 1,000 sq. ft. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS - For control of Flea Beetles, Harlequin Bugs, apply approximately $\frac{3}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For Corn Earworm, Stink Bug, Armyworm, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEAS AND OTHER ORNAMENTALS - For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillar, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

DOGS, CATS - For control of Fleas, Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens or puppies under 4 weeks old. Do not use on pregnant dogs.

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. DO NOT take internally. Wear regular long-sleeved work clothing. Change to clean clothing daily. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

1107

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

SEVIN® is a registered trademark of Rhône Poulenc Ag Company.

Label of Record - as submitted under 5/21/92 Notification

DRAGON[®]
BLUE DRAGON[®] GARDEN DUST

EPA Reg. No. 16-12
EPA Est. 16-VA-1

KILLS: Japanese Beetles, Mexican Bean Beetles, Flea Beetles, Tomato Hornworm, Corn Earworm and certain other listed insects

Contains SEVIN[®] brand carbaryl insecticide

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	2%
INERT INGREDIENTS:	<u>98%</u>
TOTAL:	100%

Keep Out of Reach of Children
CAUTION

NET WEIGHT 4 LB

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. **DO NOT** take internally. Wear regular long-sleeved work clothing. Change to clean clothing daily.
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, areas where surface water is present, or areas below the mean high tidal mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from feed and foodstuffs.
PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

GENERAL INFORMATION

BLUE DRAGON GARDEN DUST controls certain listed insects which attack vegetables, flowers, shrubs and trees. Apply when insects first appear before damage from feeding occurs. Dust thoroughly to upper and lower leaf surfaces, if needed, repeat 7-10 days. Do not use with lime or other alkaline materials. Blue Dragon may be applied up to harvest except where noted. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

STAINING PRECAUTION: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics and carpet. Deposits on painted or stained surfaces or finishes (e.g., cars, house trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid application to surfaces where visible residues are objectionable. Staining of fabrics and carpet may be permanent.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

BEANS - For control of Mexican Bean Beetle, Leafhopper, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillar, apply approximately 1½ lb. per 1,000 sq. ft. For control of Lygus, Stink Bugs, Corn Earworm, Tarnish Plant Bug, apply approximately 1½ lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN - Use approximately 1½ lb. to 2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap beetles, leafhoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS - For control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, apply approximately $\frac{1}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs, apply approximately $1\frac{1}{4}$ lb. per 1,000 sq. ft.

CUCUMBERS, SQUASH, PUMPKINS, MELONS (Cantaloupe, Watermelon, Honeydew) - For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs, Flea Beetles, Leafhoppers, apply approximately $\frac{1}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft.

LETTUCE, COLLARDS, KALE, TURNIPGREENS, MUSTARDGREENS, SPINACH - For control of Flea Beetles, Leafhoppers, Harlequin Bug, apply approximately $\frac{1}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For six-spotted Leafhoppers, Corn Earworm, Armyworm and Stink Bugs, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 14 days of harvest (except head lettuce - 3 days).

ASPARAGUS - For control of Asparagus Beetle on spears, apply approximately $1\frac{1}{4}$ to $2\frac{1}{4}$ lbs. per 1,000 sq. ft. Do not repeat application within 3 days. Do not apply within 1 day of harvest.

CABBAGE, CARROTS, BROCCOLI, BRUSSELS SPROUTS - For control of Flea Beetles, Harlequin Bugs, apply approximately $\frac{1}{4}$ to $1\frac{1}{4}$ lb. per 1,000 sq. ft. For Corn Earworm, Stink Bug, Armyworm, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest.

ROSES, AZALEAS AND OTHER ORNAMENTALS - For control of Japanese Beetle, Bag Worm, Flea Beetles, Leafhoppers, Scale Insects, Tent Caterpillar, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

GRAPES - For control of Leafhoppers, Japanese Beetle, Grape Berry Moth, June Beetles, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft.

STRAWBERRIES - For control of Meadow Spittle Bug, Leaf Roller, Strawberry Weevil, apply approximately $1\frac{1}{4}$ to 2 lbs. per 1,000 sq. ft. Do not apply within 1 day of harvest.

DOGS, CATS - For control of Fleas, Brown Dog Ticks, apply liberally and rub into skin, feet and legs. Dust sleeping quarters and infested areas weekly. Do not treat kittens or puppies under 4 weeks old. Do not use on pregnant dogs.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS and CAUTIONS stated on this label.

Manufactured by
DRAGON CORPORATION
Roanoke, VA 24019

SEVIN[®] is a registered trademark of Rhône Poulenc Ag Company.